

Half a century of *UBV* photometry at Hvar

I. Overview and emission-line stars [★]

H. Božić¹, P. Harmanec², J. Horn^{★★3}, K. Juza^{***3}, K. Pavlovski⁴, P. Koubský³, S. Kříž^{****3}, P. Mayer^{†2}, P. Hadrava³,
D. Ruždjak¹, D. Sudar¹, S. Štefl^{‡3}, M. Wolf², P. Zasche², M. Brož², J. Havelka^{§3}, J. Honsa³, V. Kocourek³,
M. Tlamicha³, F. Žďárský³, A. Harmanec⁵, J. Jonák², J. Nemravová², A. Oplištilová², I. Skokić¹, J. Švrčková²,
K. Vitovský², D. Vršnak⁶, and M. Zummer⁵

¹ Hvar Observatory, Faculty of Geodesy, Zagreb University, Kačićeva 26, HR-10000 Zagreb, Croatia

² Astronomical Institute, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, CZ-180 00 Praha 8, Czech Republic

³ Astronomical Institute, Academy of Sciences, Fričova 298, CZ-251 65 Ondřejov, Czech Republic

⁴ Department of Physics, Faculty of Science, University of Zagreb, HR-10000 Zagreb, Croatia

⁵ Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic

⁶ Faculty of Electrical Engineering and Computing, University of Zagreb, Unska 3, HR-10000 Zagreb, Croatia

Received November 12, 2024

ABSTRACT

We present a summary report on 50 years of systematic photoelectric *UBV* observations of early-type emission-line stars secured by a number of mainly Croatian and Czech observers at the Hvar Observatory. Observations of various binaries, Ap stars and several other variables will be reported in a follow-up paper. All these observations were reduced to the standard *UBV* system defined by the Johnson standard stars via non-linear transformation formulae. Our observations indicate that all Be stars are light and colour variables, but there is a great individuality in the behaviour of each studied star. We attempted to classify the variability patterns only on the basis of Hvar photometry, complemented by the Hipparcos H_p photometry transformed to Johnson *V* passband, since these observations conveniently fill-in the gap in Hvar observations caused by the war in former Yugoslavia. We still plan future detailed studies of individual well observed objects, which will be based on all their observations available in the literature and/or data archives. An important message of our effort is that to understand the complex nature of Be stars and their varying circumstellar matter, one needs to take into account light and colour behaviour in several standard photometric passbands. To rely on – accurate, but very broad band – space photometries only is somewhat dangerous since the differences in the time behaviour in the ultraviolet, blue, and yellow parts of the spectrum are inevitably washed out in those observations.

Key words. stars: early-type – stars: emission-line (Be) – binaries: eclipsing – binaries: ellipsoidal

1. Introduction

The astronomical observatory at Hvar, Croatia was founded in 1972 as a joint enterprise of several astronomical institutions from former Yugoslavia and Czechoslovakia. Two principal instruments were a solar patrol telescope and a 0.65 m Cassegrain reflector with a single channel photoelectric photometer operating with the *UBV* filters. The early history of the observatory was summarised by Mayer (2013) and Harmanec & Božić (2013) and the instrumentation and the original reduction techniques were described in Mayer (1977) and Harmanec et al. (1977a).

First observations were secured on July 29, 1972 and continued more or less systematically since then, with the only longer gap (caused by the war in the disintegrating Yugoslavia) from February 1991 until September 1993.

During the first, 1972 observing season, we primarily tried to cover a complete light curve of the short-periodic eclipsing binary BR Cyg to demonstrate the good performance of the new instrument and this result was soon published (Harmanec et al. 1973a). In January 1973, Pavel Mayer obtained a nearly complete light curve of another eclipsing binary, IU Aur, and confirmed his 1971 discovery that the depths of the light-curve minima of this system are secularly changing due to precession of the binary orbit caused by the presence of a third body in the system (Mayer 1976).

Our main motivation at that time was, however, an attempt to test the binary hypothesis of the origin of the Be phenomenon (Kříž & Harmanec 1975; Harmanec & Kříž 1976). The idea was that if Be stars are indeed mass accreting components of binary system in the stage of large-scale mass transfer, with Roche-lobe filling secondaries, one should find a non-negligible number of eclipsing and ellipsoidal binaries among them. While we later indeed found several such systems, for instance CX Dra (Koubský et al. 1980; Horn et al. 1992; Koubský et al. 1998), KX And (Harmanec et al. 1977b; Štefl et al. 1990), V360 Lac (Hill et al. 1997; Linnell et al. 2006), BR CMi (Harmanec et al. 2015), or HD 81357 (Koubský et al. 2019), for the majority

[★] Based on photometric observations from the Hvar Observatory and ESA Hipparcos Satellite.

^{★★} Jiří Horn passed away on Dec. 13, 1994

^{***} Karel Juza passed away on March 13, 1994

^{****} Svatopluk Kříž passed away on Feb. 23, 2018

[†] Pavel Mayer passed away on Nov. 7, 2018

[‡] Stanislav Štefl passed away on June 11, 2014

[§] Josef Havelka passed away on June 19, 2009

Table 1: A complete list of Hvar observers and the number of usable individual differential observations per observer, which they secured at the Hvar Observatory over the years 1972 - 2024

Name of observer	No. of observations
Hrvoje Božić	46237
Petr Harmanec	9530
Krešimir Pavlovski	8674
Domagoj Ruždjak	5636
Davor Sudar	5259
Jana Nemravová	3945
Vaclav Kocourek	3876
Miloslav Tlamicha	3715
Petr Hadrava	3485
Alžbeta Oplištilová	2859
František Žďárský	2752
Kristian Vitovský	2064
Jiří Horn	1927
Pavel Koubský	1924
Jaroslav Honsa	1902
Petr Zasche	1849
Miroslav Brož	1837
Svatopluk Kříž	1805
Marek Wolf	1722
Josef Havelka	1607
Jana Švrčková	1552
Michal Zummer	1474
Damir Čikotić	1449
Donik Vršnak	1436
Rostislav Brož	1250
Juraj Jonák	1036
Jiří Grygar	988
Stanislav Štefl	964
Karel Juza	626
Adam Harmanec	606
Isabelle Piantschitsch	582
Ivica Skokić	515
Jakub Juryšek	417
Mirjana Malarić	373
Pavla Polechová	347
Miloš Zejda	334
Jan Janík	334
Pavel Chadima	327
Zoran Ivanović	315
David Ondřích	302
Željko Ružić	269
Dunja Plačko-Vršnak	218
Istvan Vince	209
Pavel Mayer	177
Kateřina Hoňková	166
Gordana Lazin	137
Jiří Vacek	136
Muhamed Muminović	136
Martin Netopil	121
Emil Frlež	99
Jelisaveta Arsenijević	83
M. Mulić	68
Boro Jovanović	43
Roman Brajša	24
Aldo Arena	24
Bojan Vršnak	17
Željko Ivezić	6

of the observed Be stars we discovered long-term light and colour changes related to the changes in their circumstellar envelopes (seen in parallel spectroscopy), on a time scale of years; see, for instance Harmanec et al. (1976); Doazan et al. (1982b,a); Horn et al. (1982) and the three summary reports (Harmanec et al. 1980b; Pavlovski et al. 1997; Harmanec et al. 1997).

The first three selected Be stars, V744 Her = 88 Her, V439 Her = 4 Her, and EW Lac = HD 217050 were observed already in 1972. Our discovery that essentially all Be stars are light and colour variables was quite important. At that time, Be stars were sometimes used as the comparison stars in various observing programs of differential photometry. Perhaps the most interesting is the case of V1294 Aql = HD 184279, which was recommended as the secondary standard of the *UBV* system – see this story in Horn et al. (1982). That prompted us to prepare a list of all at that time known bright Be stars and select for each of them suitable comparison and check stars. Since 1980 we organised an international observing campaign, which was coordinated mainly via the Be Star Newsletter, created and for a long time run by M. Jaschek (Harmanec 1980; Harmanec et al. 1980a, 1982). We provided the list of Be stars with the recommended comparisons to interested observers and reported their observations in the Be Star Newsletter.

Around 1980, first studies appeared, in which low-amplitude periodic or even multiperiodic light variations of some Be stars were detected (Bolton 1982; Percy 1982, 1983; Harmanec 1984b; Balona & Engelbrecht 1986). Since the typical timescales of such variations were from about 0.3 to 2 d, observations secured at any single observing station suffered from strong aliasing problems. This led us to organise a collaborative observing campaign with the participation of several observers located in very different geographic longitudes all around the globe. The campaign was carried out in 1983 and was quite successful. Preliminary results were reported by Stagg et al. (1985) and the final report was published three years later (Stagg et al. 1988).

Since 1973, there was also a program of photometric monitoring of selected chemically peculiar (CP) stars, both single and binary ones. The first result of this effort were two detailed studies of the oblique rotator model of the CP star CQ UMa (Mikulášek et al. 1978; Pavlovski 1979).

Later on, also monitoring of various, mainly eclipsing binaries has also been on the program of the Hvar photometer.

A complete list of observers who participated in some observations at Hvar and the number of usable differential observations they secured is in Table 1.

2. The observational strategy and the new reduction of all observations since 1972

Since our motivation, especially for the dominant Be-star program, was to ensure stable and as accurate as possible transformation to the standard Johnson system, we soon found that we have to use non-linear transformation formulae for the conversion of extinction-free instrumental magnitudes to the standard Johnson magnitudes, and to derive the coefficients of these transformations from all observations of standard stars in each given observing season.

Originally, our reduction program used a linear extinction coefficient as a function of the air mass, which was assumed to remain constant during each observing night. This is actually almost never so, even at observing stations with good climate conditions like La Silla or Sutherland. For Hvar Observatory, located

Table 2: The robust mean values of the *V* magnitude and the *B* – *V*, *U* – *B* and *V* – *R* indices for the ten primary Johnson standards derived from individual observations published by Johnson et al. (1966).

Name	HD	<i>V</i>	rms	<i>N</i>	<i>B</i> – <i>V</i>	rms	<i>N</i>	<i>U</i> – <i>B</i>	rms	<i>N</i>	<i>V</i> – <i>R</i>	rms	<i>N</i>
α Ari	12929	2.0060	0.0033	191	1.1509	0.0012	192	1.1260	0.0018	189	0.8432	0.0039	169
HR 875	18331	4.2836	0.0020	77	-0.0551	0.0018	78	-0.1165	0.0021	63	0.0160	0.0029	63
β Cnc	60267	3.5231	0.0019	223	1.4830	0.0067	225	1.7699	0.0020	224	1.1155	0.0021	197
η Hya	74280	4.2985	0.0012	186	-0.1941	0.0011	186	-0.7446	0.0014	182	-0.0634	0.0021	157
β Lib	135742	2.6072	0.0017	143	-0.1073	0.0015	143	-0.3614	0.0017	143	-0.0384	0.0040	128
α Ser	140573	2.6355	0.0014	135	1.1704	0.0014	137	1.2509	0.0021	137	0.8109	0.0048	123
ϵ CrB	143107	4.1527	0.0032	127	1.2324	0.0012	126	1.2751	0.0025	126	0.9012	0.0036	114
τ Her	147394	3.9052	0.0020	102	-0.1522	0.0019	106	-0.5621	0.0018	103	-0.0822	0.0034	88
10 Lac	214680	4.8845	0.0014	204	-0.2045	0.0012	204	-1.0464	0.0014	203	-0.0751	0.0018	187
HR 8832	219134	5.5738	0.0019	158	0.9966	0.0014	158	0.8861	0.0018	158	0.8346	0.0023	144

Table 3: The robust mean values of the *V* magnitude and *B* – *V*, *U* – *B* and *V* – *R* indices derived from individual observations of seven selected secondary standards (Johnson & Harris 1954) by Johnson et al. (1966).

Name	HD	<i>V</i>	rms	<i>N</i>	<i>B</i> – <i>V</i>	rms	<i>N</i>	<i>U</i> – <i>B</i>	rms	<i>N</i>	<i>V</i> – <i>R</i>	rms	<i>N</i>
73 Cet	15318	4.2836	0.0020	77	-0.0551	0.0018	78	-0.1165	0.0021	77	0.0160	0.0029	63
HR 753	16160	5.8164	0.0077	10	0.9633	0.0039	10	0.8074	0.0078	10	0.8177	0.0095	7
ν Ori	36512	4.6215	0.0014	273	-0.2618	0.0099	276	-1.0626	0.0012	274	-0.1217	0.0015	246
90 Leo	100600	5.9495	0.0014	170	-0.1547	0.0011	170	-0.6572	0.0013	168	-0.0589	0.0026	129
HR 4550	103095	6.4411	0.0017	167	0.7511	0.0012	169	0.1708	0.0016	167	0.6637	0.0020	131
β Oph	161096	2.7619	0.0023	105	1.1689	0.0018	107	1.2408	0.0018	106	0.8138	0.0033	88
γ Oph	161868	3.7471	0.0024	110	0.0380	0.0015	111	0.0348	0.0021	110	0.0457	0.0042	91

on an island at an altitude of 260 m only above the sea level, the variations of the linear extinction during the night are usually quite large, especially during summer seasons. To cope with this problem, we had to split many observing nights to shorter segments in time and to derive extinction for each segment separately. This was not satisfactory since the standard stars from many - otherwise good and long observing nights - could not be used for seasonal transformations.

Only much later a version of the reduction program was developed, which allows the monitoring of the variable extinction during the observing night. This was one of the reasons, why we decided to undertake the tedious task to re-reduce all observing seasons since 1972.

The other one was the following: The strategy of Hvar observations, developed first for the Be star program but later used for all observing programs, was that we prepared a list of all known bright Be stars north of declination of -30° , which were listed in the Bright Star Catalogue (Hoffleit & Jaschek 1982), and a few fainter ones. For all of them, we selected suitable comparison and check stars. If more Be stars were found close to each other in the sky, we observed them altogether, using the same comparison, and check star. Since most of our observations dealt with early type, blue stars, we also selected the so called red standards for each group to facilitate good seasonal transformation. Whenever possible, we have chosen the comparisons, for which the *UBV* photometry obtained by Johnson and his collaborators was available, our primary source being the summary publication by Johnson et al. (1966). It was, of course, a must for the red standards. This allowed us to save the observing time. Com-

parisons, checks, and red standard stars with known *UBV* magnitudes could be used to both, the monitoring the extinction during observing nights, and to seasonal transformation to the standard *UBV* system.

It turned out, however, that the stars, which have only a few original Johnson observations suffered from rather large accidental errors. Our task then was to improve the original Johnson values but to preserve the original Johnson system over the whole range of the Hertzsprung-Russel (HR) diagram.

Several years ago, we decided to derive robust mean *UBVR* values for all individual observations of the recommended Johnson primary standards, for which extinction was measured, from Johnson et al. (1966) study and use them in all consecutive observations at Hvar during good nights, suitable for all-sky photometry. These are provided in Table 2. Additionally, we did the same also for several well observed stars from the list of 108 recommended Johnson secondary standards (Johnson & Harris 1954). Their robust *UBVR* values are listed in Table 3. All selected stars were carefully checked for variability with the help of individual H_p space-born Hipparcos photometry (Perryman & ESA 1997) and SIMBAD bibliography¹. Of the ten primary Johnson standards two are reported as variables. η Hya is classified as the β Cep pulsator and τ Her as a long-period pulsating star. α Ari, β Cnc, β Lib; α Ser, 10 Lac, and HR 8832 are listed as suspected variables. Of the seven selected secondary standards, β Oph is listed as a suspected vari-

¹ <http://simbad.u-strasbg.fr/simbad/sim-fid>

able. HR 4550 got a variable-star name CF UMa and deserves a more detailed comment (to be done later).

When analysing the errors of the Hipparcos H_p observations, as well as the errors of the Hvar photometry of Johnson standards used, we conclude that these errors usually do not exceed 0^m.01. As shown already by Harmanec et al. (1994), the usual scatter of individual observations for non-variable stars at Hvar is within 0^m.02 to 0^m.03. Accordingly, the use of the selected Johnson standards, even those having possible low-amplitude variations, has no systematic effect on the stability of the Hvar standard photometric system.

The Johnson standards were then systematically observed several times during each good observing night and a care was taken to observe them in high and low air masses to bracket observations of all other stars during each observing night. We also carried out dedicated observations of all important comparison and check stars, for which good Johnson values were missing. This effort continued for several last years and allowed us to get better truly Johnson values for constant stars used in the early observing seasons. With them, we repeated the reductions of all seasons since 1972. In the vast majority of cases this resulted in much better seasonal transformations and a significant decrease of the mean standard errors.

In particular, we used the following working equations.

For each star and the time of observation, we derived the zenith distance z from its equatorial coordinates recalculated for the time of observation and derived the accurate air mass after Bemporad (1904):

$$X = \sec z - 0,0018167Q - 0,02875Q^2 - 0,0008083Q^3, \quad (1)$$

where

$$Q = \sec z - 1. \quad (2)$$

The transformation coefficients G between the observed instrumental magnitudes, denoted v , b , u , and r , and extinction-free magnitudes v_0 , b_0 , u_0 , and r_0 , determined (or fixed) separately for each night of observations include the first-order extinction coefficients and a possibility to model time variations of the linear extinction coefficients during each observing night via polynomials up to 5th degree as a function of time t . The corresponding transformation equations read as

$$\begin{aligned} v &= v_0 + G_1 + G_5 X + G_9 tX + G_{13} t^2 X \\ &+ G_{17} t^3 X + G_{21} t^4 X + G_{25} t^5 X, \\ b &= b_0 + G_2 + G_6 X + G_{10} tX + G_{14} t^2 X \\ &+ G_{18} t^3 X + G_{22} t^4 X + G_{26} t^5 X, \\ u &= u_0 + G_3 + G_7 X + G_{11} tX + G_{15} t^2 X \\ &+ G_{19} t^3 X + G_{23} t^4 X + G_{27} t^5 X, \\ r &= r_0 + G_4 + G_8 X + G_{12} tX + G_{16} t^2 X \\ &+ G_{20} t^3 X + G_{24} t^4 X + G_{28} t^5 X. \end{aligned} \quad (3)$$

There is also a possibility to take into account a linear or quadratic drift of the zero point of the instrument. In that case, the drift is derived via the yellow-magnitude (V or y) observations and forced to data in all other passbands. The corresponding transformation equations then read as

$$\begin{aligned} v &= v_0 + G_1 + G_5 X + G_9 t + G_{13} t^2, \\ b &= b_0 + G_2 + G_6 X + G_9 t + G_{13} t^2, \\ u &= u_0 + G_3 + G_7 X + G_9 t + G_{13} t^2, \\ w &= w_0 + G_4 + G_8 X + G_9 t + G_{13} t^2. \end{aligned} \quad (3)$$

The colour transformation equations between the Johnson standard value for each magnitude (V, B, U, R) and the extinction-free instrumental magnitudes are linear in the $U - B$ index but have the form of a third-degree polynomial in $B - V$. This form of the equations is necessary to compensate for the unavoidably non-linear effect of the Balmer jump on the magnitudes of stars between mid-B to F spectral types (see Cousins & Jones 1976). The colour extinction coefficients in the form recommended by Young et al. (1991) are also included among the seasonal transformation coefficients. It is our experience that this form of transformation equations ensures the reproduction of the standard Johnson system within 0^m.01 even in U for any standard star.

The equations are:

$$\begin{aligned} v_0 - V &= H_1(B - V) + H_2(U - B) + H_3(B - V)^2 \\ &+ H_4(B - V)^3 + H_5 X B_4(B - V + 0.5 X B_4) + H_6, \\ b_0 - B &= H_7(B - V) + H_8(U - B) + H_9(B - V)^2 \\ &+ H_{10}(B - V)^3 + H_{11} X B_4(B - V + 0.5 X B_4) + H_{12}, \\ u_0 - U &= H_{13}(B - V) + H_{14}(U - B) + H_{15}(B - V)^2 \\ &+ H_{16}(B - V)^3 + H_{17} X B_5(U - B + 0.5 X B_5) + H_{18}, \\ r_0 - R &= H_{19}(B - V) + H_{20}(U - B) + H_{21}(B - V)^2 \\ &+ H_{22}(B - V)^3 + H_{23} X B_5(U - B + 0.5 X B_5) + H_{24}, \end{aligned}$$

where $B_4 = G_6 - G_5$ and $B_5 = G_7 - G_6$ are the linear extinction coefficients in the $B - V$, and $U - B$ colours.

Please note that in this formalism, quantities with suffix 0 are not in fact really extinction-free instrumental magnitudes but extinction-free magnitudes uncorrected for the colour extinction which is only taken into account in the seasonal transformation equations, where they actually belong since they depend on the colour properties of the photometer, not on the properties of the Earth atmosphere.

We warn that it is necessary to obtain observations of a reasonably large number of standard stars of different colours, luminosity classes and reddenings to derive reliable non-linear transformation coefficients H . Since this is not always possible, e.g., due to bad weather, the latest versions of the program also permit the use of bilinear transformation formulae which contain no terms of the second and third power in $(B - V)$ and converge to reasonable transformation coefficients in most situations. This compromise is, however, at the expense of a slightly worse transformation into the standard system, yet still much better than would result from the use of the often applied linear transformation formulae. The possibility to derive and use the linear transformation formulae is also included (non-zero coefficients H_1 and H_6 , H_7 and H_{12} , H_{13} and H_{18} , and H_{19} and H_{24}). This can be justified in situations when the observations only include objects with a limited range of colours or for passbands covering the flat parts of the stellar energy distributions (R or even V , for instance).

Since the non-linear transformations we used imply that all three passbands affect the accuracy of the determination of the standard values, we took care for each observed star to use different integration times, usually longer for the U passband, to obtain comparable signal-to-noise (S/N) ratio for all of them.

3. Long-term, orbital, and rapid light and colour changes of Be stars observed at Hvar

The list of all Be stars observed at Hvar is in Table 6, while more detailed information about the observations, time interval covered by observations, and the range of variations, is summarised in Table 7.

Our long experience with the photometry of Be stars as well as numerous studies of other colleagues all over the world allowed us to recognize the following typical patterns of light and colour changes:

- Long-term changes related to the strength and/or extent of the Be envelope (LTEM). These occur on a time scale of years to decades and have two distinct alternatives, depending on the inclination of the Be disk plane with respect to the observer: positive (LTEM_p), when the brightenings are accompanied by the rise of the emission-line strength, and inverse (LTEM_i), when the light decreases are followed by the emission-line rise (Harmanec 1983, 1994, 2000; Sigut & Patel 2013). We note that in cases of intermediate inclination, given object can exhibit either a positive, or an inverse correlation in various epochs of observations, depending on the extent of the Be envelope (see, e.g. Harmanec et al. 2022).
- Cyclic long-term changes related to the long-term cyclic *V/R* variations of the relative strength of the *V* and *R* peaks of the double Balmer emission lines (LTVR). Such light changes occur typically on a time scale of several years and are related to non-axisymmetric geometrical and/or dynamical changes of the structure of the Be disks. In the past, they were interpreted as a gradual revolution of an elongated disk (see, e.g. Ballereau & Chauville 1989). Since the study by Okazaki (1991), they are usually understood as global one-armed oscillations of the disks and can also manifest themselves by corresponding light changes (see the detailed discussion by Mennickent et al. 1997). A recent example of such changes was presented by Wolf et al. (2021). The trouble is that the LTVR occur sometimes simultaneously with the LTEM changes and it is not easy to separate both effects (see, for instance Harmanec et al. 2022).
- Light and colour variations related to the duplicity of particular Be stars (MTBIN). Such changes are periodic and generally follow the orbital periods of the Be binaries in question but they are not easy to be detected due to the simultaneous presence of variations on other time scales. Besides clear cases of binary eclipses, they can also manifest themselves as a combination of ellipticity and reflection or occasional eclipses of the outer parts of the disks. Numerous examples of such changes were discussed, with corresponding references, by Božić et al. (2013a) and are documented by the Hvar systematic photometry – see the discussion of individual Be stars below.
- Rapid periodic or multiperiodic low amplitude changes (RAV). These brightness variations usually occur on timescales from about 0^d.1 to 2 d and have been alternatively interpreted as pulsations or rotational modulation. Since the typical observing strategy at Hvar was focused on monitoring of variations on longer time scales, Hvar data are not very suitable for analyses of such rapid changes, with the exception of several dedicated observing campaigns on selected Be stars, for instance ζ Tau (Božić & Pavlovski 1988), σ And, KX And, KY And, LQ And and EW Lac (Stagg et al. 1988) or ω Ori (Balona et al. 2001).
- Secular very long-term brightness decreases or increases during quiescent stages without strong LTEM changes (LTCV). This very long, “fourth” timescale of changes was probably first noted for ω CMa by Harmanec (1998b) and modelled empirically in terms of

the viscous decretion disk model by Ghoreyshi et al. (2021, 2023). Several distinct examples of such behaviour were presented by Harmanec et al. (2022). The understanding of this type of variability will require further observational and theoretical effort.

Below, we discuss particular stars, which were observed more systematically at Hvar. Because the observations at Hvar were interrupted during 1990-1992, at the times of the war in Yugoslavia, we complemented the Hvar *V* series in the *V* magnitude time plots with the Hipparcos *H_p* photometry, transformed to Johnson *V* magnitude after Harmanec (1998a). These are shown by red circles in all plots. The following notes about particular observed Be stars are mostly related to LTVR, LTEM, LTCV, and MTBIN changes.

σ Cas = 22 Cas = HD 4180 This bright Be star is the primary component A of a wide visual system AB WDS J00447+4817. Koubský et al. (2010) confirmed that it is also a spectroscopic binary with a period of 1031^d.6 and a rather large RV semi-amplitude of 22 km s⁻¹, which implies a large mass function. They also obtained a visual orbit with an inclination of 115° and since they did not detect any lines of the secondary, they suggested that the secondary is actually a close pair of two A stars. Such stars were indeed detected in the spectra studied by Grundstrom (2007), with a possible period close to 4 days (see also Touhami et al. 2013; Hutter et al. 2021). It thus appears that the object is a hierarchical system Aa1, Aa2, Ab1, Ab2, and B. Koubský et al. (2010) also found a period of 1^d.2578 in the residuals from their RV orbit and showed large-amplitude cyclic light changes, based on the Hvar photometry. Here, we present extended series of Hvar observations in Fig. 1.

γ Cas = 27 Cas = HD 5394 This one of the brightest Be stars in the sky is a subject of more than 1100 studies. It is now known to be a spectroscopic binary with a 203^d.52 orbital period (Harmanec et al. 2000; Nemravová et al. 2012; Borre et al. 2020, and references therein), with a low mass companion, probably a white dwarf (Gies et al. 2023). Its light variations are characterised by long-term changes with a positive correlation and rapid changes with periods 1^d.216, and 0^d.4033 (Harmanec 2002; Borre et al. 2020; Smith & Henry 2021, and references therein). Observations of this star at Hvar are not very numerous, mainly because of the fact that it was necessary to use a lower high voltage due to its brightness to avoid saturation. They are shown in Fig. 2 and only complement the secular brightness changes documented in Harmanec (2002) and Smith & Henry (2021).

V442 And = HD 6226 This star was originally used as one of the check stars in our Be star observing program and this led to the discovery of its light variability in the form of occasional brightenings, accompanied by the reddening in *B*–*V*, and blueing in the *U*–*B* index. This finding was published by Božić & Harmanec (1998), who suggested that the object could be a Be star. This was indeed confirmed by a detailed spectroscopic and photometric study by Božić et al. (2004), who found that V442 And is a pole-on Be star with a positive correlation between brightness and emission-line strength and pronounced line-profile variations, reminiscent of an archetype Be star ω CMa, and varying with a regular 2^d.61507 period. Richardson et al. (2021) published a very detailed study, based on numerous spectra and on the KELT and TESS photometries. They estimated the spectral type as B2.5IIIe and confirmed the




Fig. 1: The *UBV* time variations of *o* Cas. The red dots in this, and in all following time plots for individual objects denote the Hipparcos H_p observations transformed to Johnson *V* magnitude after Harmanec (1998a).




Fig. 2: The *UBV* time variations of γ Cas.

period of line-profile changes. Analysing the $H\alpha$ and $H\beta$ equivalent widths, they found regular changes with two periods, $211^{\text{d}}9$, and $87^{\text{d}}0$. They also argued that the emission-line episodes could be related to the interference of pulsational modes, resulting in formations of transient circumstellar disks. Other published studies are also summarised in their paper.

The complete light curve of V442 And secured at Hvar is in Fig. 3. There is some suspicion that the ‘undisturbed’ brightness of the object outside the brightenings might be decreasing secularly on the fourth time scale.

φ And = 42 And = HD 6811 This Be star is obviously observed pole-on, shows a single-peaked $H\alpha$ emission profile and little light variability, and was alternatively classified as B7Ve and B5IIIe (Jones et al. 2011; Barnsley & Steele 2013). However, the available records of its $H\alpha$ line profile show large secular changes up to almost complete disappearance of the emission - see, e.g. the $H\alpha$ spectra available in the BeSS database (Neiner et al. 2011). It is also a member of a multiple visual system WDS J01095+4715 and Muterspaugh et al. (2010) published a visual orbit with a period of about 202500 d. Its mild light and colour variability was discovered at Hvar thanks to a very systematic monitoring and careful reduction – see Fig. 4. It is characterised by a slow light decrease on the fourth time scale, with no known episodes of more rapid brightening or fading.

φ Per = 54 And = HD 10516 This object is a well-known B0.5e+O6 VI binary with a $126^{\text{d}}7$ period (see Poeckert 1981a,b; Gies et al. 1993; Thaller et al. 1995; Božić et al. 1999; Gies et al. 1998, and references therein). Božić et al. (1995) compiled all at that time available photometric and RV observations and summarised the early history of the system. They







Fig. 3: The *UBV* time variations of V442 And.

also derived an improved orbital period of the system. Photometry, including the first part of Hvar observations, shows only mild secular changes. They interpreted the emission seen in $\text{He I } 6678 \text{ \AA}$ as a double emission moving with the secondary. However, Štefl et al. (2000) and Hummel & Štefl (2001) argued


Fig. 4: The *UBV* time variations of ϕ And.

Fig. 6: The *UBV* time variations of V554 Per.

Fig. 5: The *UBV* time variations of ϕ Per.

Božić et al. (2013b) reported small sinusoidal *B* magnitude light variation with the orbital period.

Complete Hvar photometry is shown in Fig. 5. It is seen that the light variations are dominated by slow changes on the fourth time scale. The *V* magnitude varies in a cycle of more than 5000 d, while the *B*–*V* index was gradually brightening over many years but started to get redder recently.

V554 Per = 10 Per = HD 14818 This object is classified B2Ia but Weißmayer et al. (2022) observed its H α emission, reminiscent of a P Cyg profile. Its light variability was reported by Alvarez & Schuster (1981) and Pavlovski et al. (1997) but the variable-star name V554 Per was assigned to it only on the basis of Hipparcos photometry by Kazarovets et al. (1999).

Hvar and Hipparcos photometry is shown in Fig. 6. The brightness and colour variations are characterized by cyclic changes, with no obvious secular trend. A formal period analysis of combined Hvar and Hipparcos *V* photometry suggests a possible period of 237^d082, but with a large scatter around the mean phase curve.

HR 894 = HD 18552 Polidan (1976) tentatively suggested that this object is a B8V+gG9 binary, based on the calcium triplet emission in the infrared spectra. However, the presence of a cool giant companion has never been confirmed. The object exhibits only mild variations of the H α emission over a long interval of spectral observations and Hvar photometry shows no significant light or colour changes (see Fig. 7). We thus did not continue observing it.

RX Cas = BD+67 244 This object is an interacting Be+G3III binary with a 32^d3 period secularly increasing in time and with a cyclic brightness variation at light maxima on a scale of about 500 days (Struve 1944; Gaposchkin 1944; Kříž et al.

that this emission actually originates in the part of the disk of the primary, facing the hotter secondary and being illuminated by it. Mourard et al. (2015) resolved the secondary in their optical interferometry and published the so far most detailed model of the system, deriving the component masses of 9.6 and 1.2 M_{\odot} .




Fig. 7: The *UBV* time variations of HR 894.




Fig. 8: The *UBV* time variations of RX Cas.

1980; Andersen et al. 1989). Mennickent et al. (2022) analysed more than a century of photometric observations, derived an improved quadratic ephemeris and studied the long cycle of brightness variations at maximum light. To the best of our





Fig. 9: The magnitude and colour phase diagrams for RX Cas based on the quadratic ephemeris of Mennickent et al. (2022).

knowledge, there is no reliable RV curve of the Be primary hidden in the dense shell.

The first part of Hvar photometry has already been used in Kříž et al. (1980) study. The complete set of Hvar observations is shown in Fig. 8. In this case we have not included the Hipparcos photometry since the *B*–*V* index of the binary varies significantly with the binary orbital phase and with the long cycle and the transformation to Johnson *V* would deserve a more sophisticated treatment. Our new detailed study of the binary is under consideration and here we only show the phase coverage based on the complete set of Hvar data in Fig. 9. It is seen that the phase coverage is quite good. We note that the secondary minimum is absent in the *U* magnitude.



440 13 Tau = HD 23016 is a Be star which is only projected in the sky to the Pleiades cluster but is not its member. Barnsley & Steele (2013) provide some evidence of a mild change in the *H*α emission strength (equivalent range from 4.0 460


Fig. 10: The *UBV* time variations of 13 Tau.

to 5.7 Å) over the time interval from JD 2451029 to 2455524. Hvar photometry shown in Fig. 10 provides little evidence for a significant light variability over a time interval of nearly 10000 days.



17 Tau = HD 23302 This bright member of the Pleiades cluster, also known as Electra, was reported to be a single-line spectroscopic binary with the orbital period of 100^d46 by Abt et al. (1965). They claimed that the orbit, based on a limited set of RV measurements, is not well determined. Pearce & Hill (1975) were not able to definitely provide proof that the star was spectroscopic binary but left the possibility open. Torres (2020) derived accurate RVs over an interval of about 800 days and concluded that there is no evidence of RV changes. Breger (1972) showed that 17 Tau is constant down to 0^m002-0^m003. In his search for Maia variables McNamara (1985) used 17 Tau as a comparison star in the photometric search for the Maia candidates, finding it to be constant, and its photometric constancy was also confirmed by McNamara (1987). Also the Hipparcos H_p photometry does not show any significant brightness changes. The Hvar photometry presented in Fig. 11 shows the presence of mild long-term variations in brightness and colours on the level of several hundredths of a magnitude.

V971 Tau = 23 Tau = HD 23480 For this bright Be star from the Pleiades cluster, also known as Merope, (McNamara 1985, 1987) found rapid variation with the 0^d490 period and an amplitude of several thousandths of magnitude. White et al. (2017) using data from the Kepler mission confirmed that periodicity and suggested the existence of several closely spaced periods. In spite of a very small amplitude of rapid changes, the Hvar photometry is able to resolve the periodic variations with the 0^d490 period. The Hvar photometry presented in Fig. 12 indicates that the small light variability is dominated by these rapid changes


Fig. 11: The *UBV* time variations of 17 Tau.

Fig. 12: The *UBV* time variations of V971 Tau.

but reveals also mild long-term variations, best seen in the $U-B$ index.

490 η Tau = 25 Tau = HD 23630 This brightest Be member of the Pleiades cluster, also known as Alcyone, is a visual multiple system WDS 03475+2406. Figure 13 shows that mild bright-

Fig. 13: The UBV time variations of η Tau.Fig. 14: The UBV time variations of BU Tau.

ness and colour variations on various time scales are probably present. A cyclic secular variation is best seen in the $U-B$ index. It is worth mentioning that Krelowski et al. (2019) studied the distances to the brightest stars in the Pleiades derived both from Gaia DR2 release and from interstellar Ca II lines and concluded these stars do not form a compact cluster but are localized in front of the main Pleiades cluster at a distance close to what was derived from Hipparcos. They argued that the observed spread in the individual distances of the stars in the core of the cluster could be explained by additional grey interstellar extinction due to the presence of larger dust grains within in the cluster core.

BU Tau = 28 Tau = HD 23862 This well-known Be star is one of few for which brightness and colour variations have been recorded systematically for a long time, mainly thanks to the effort of Sharov and Lyutyj (see Sharov & Lyutyj 1972; Sharov & Lyutyj 1997, and references therein). It is the primary component of a binary with a highly eccentric orbit and an orbital period of 218^d03 (Nemravová et al. 2010; Katahira et al. 1996). The orbit was not resolved interferometrically by Klement et al. (2024). The spectroscopic behaviour of this star is characterised by remarkable changes between normal Be state to Be-shell and Be phase (Hirata & Kogure 1976; Hirata 1995). After a long B phase that lasted until 1937, the star passed through 3 cycles of photometric and spectral variation. The photometric variations show long-term cyclic changes which started with the light decrease lasting for some two years. After reaching the light minimum, a gradual increase in brightness to the initial level was temporarily replaced by another mild light decrease, related to the development of a metallic shell. The period of the complete cycle lasts 34-36 years (Hirata & Kogure 1976; Harmanec 1982). Using speckle interferometry McAlister et al. (1989) found a third companion at the separation of 0.22 sec. Harmanec (1982) and Gies et al. (1990) (and some others) con-




sidered a causal connection between the motion of that distant companion and the occurrence of the shell phases. Roberts et al. (2007) recorded also the fourth companion of BU Tau at a distance of 4.66 sec.

Besides a few observations from the season 1976/77, Hvar data cover almost the entire last cycle of photometric changes. A secondary minimum is visible on the ascending part of the curve, the depth of which is the most prominent at shorter wavelengths – see Fig. 14.

BU Tau shows the inverse correlation between the brightness and emission-line strength. The behaviour in $U-B/B-V$ is shown in Fig. 53.

V960 Tau = 120 Tau = HD 36576 The light variability of this star was discovered at Hvar and first reported by Pavlovski & Božić (1982). As Fig. 15 shows, the star was monitored quite systematically at Hvar and shows complicated and seemingly irregular variations with a large range of 0^m3 in V . Its detailed study, which will also include spectroscopic changes, is in preparation.

ζ Tau = 123 Tau = HD 37302 This well-known and for a long time studied Be star is a single-line spectroscopic binary with a 132^d99 orbital period, exhibiting also long-term cyclic RV and V/R changes at certain time intervals (see the detailed studies by Delplace 1970; Harmanec 1984b; Ruždjak et al. 2009; Štefl et al. 2009; Carciofi et al. 2009, where also references to original papers can be found). There is a clear correlation between the long-term RV and V/R changes. However, Ruždjak et al. (2009) have demonstrated that the light and colour behaviour was different in each long cycle, alternating between positive and inverse correlation. The first part of Hvar photometry has also been discussed by Božić & Pavlovski (1988). These authors found, in addition to long-term changes, also rapid variations with a 0^d8 period (or a 1^d6 period with a double-wave


Fig. 15: The *UBV* time variations of V960 Tau.

Fig. 17: The *UBV* time variations of ω Ori.

Fig. 16: The *UBV* time variations of ζ Tau.

phase curve). They also detected orbital modulation present in some cycles, which was missing in the others.

A complete set of Hvar observations is shown in Fig. 16. The apparent scatter of the data is mainly due to large-amplitude rapid changes.

ω Ori = 47 Ori = HD 37490 Hvar photometry was obtained mainly as a part of a multisite campaign on the rapid line-profile and light variability, which resulted in the finding that both, line-profile and light changes are governed by a single 0^d97 period (Balona et al. 2001). Several episodes of a rapid light increase have been reported (e.g. Guinan & Hayes 1984; Bergin et al. 1989) accompanied also by an increase of polarization so it is probable that this star has a positive correlation between the brightness and H α emission strength and might be similar to V442 And. Not very rich Hvar photometry can be seen in Fig. 17. It seems that the variations of this star are dominated by rapid changes, no secular changes are seen.

V731 Tau = HD 37967

A limited set of Hvar observations seems to show only small and probably rapid changes - see Fig. 18. Sigut & Ghafourian (2023) estimated the disk inclination of about 55° for this B4 star.

V696 Mon = HR 2142 = HD 41335 This Be star was found to exhibit two consecutive short-lived shell phases periodically every 80^d85 days (Peters 1971, 1972). Later, Peters (1983) found that the object is a single-line spectroscopic binary with an orbital period of 80^d860, derived its radial-velocity curve and argued that the observed shell phases are consequences of mass transfer in a binary system. Peters originally argued that the secondary is a late-type Roche-lobe filling object but no such object has ever been detected and Waters et al. (1991) proposed an alternative model, assuming that the secondary is a small helium star and that the disk around the primary is not an accretion, but an outflowing disk. Peters et al. (2016) indeed detected a weak signal in the IUE spectra corresponding to a hot subdwarf companion, with a mass ratio of 0.07 and T_{eff} around 43000 K. Using numerous optical and IUE spectra, they also revised the orbital




Fig. 18: The *UB* time variations of V731 Tau.




Fig. 19: The *UB* time variations of V696 Mon.

period to 80^d913. Sterken (1983) reported a 0^m.13 light decrease in the Strömgren *b* magnitude. Hvar photometry of the object has been secured rather systematically since the end of 1981 and it was first used, together with the Kitt Peak and Skalnaté Pleso photometry to demonstrate that no such light decrease occurred and that also the recurrent shell phases are not accompanied by any light changes (Harmanec et al. 1983). The absence of light changes during a primary shell phase was also reported by Dorren et al. (1984).

The time plot of the Hvar photometry, shown in Fig. 19 shows only a very mild secular light decrease on the fourth time scale and little or no changes in both colour indices. One could suspect some small rapid changes to be present but we note that the star is always observed at air masses larger than 1.55 at Hvar, which implies somewhat higher scatter of individual observations.

OT Gem = HD 58050 Hvar observations were already reported by Božić et al. (1982) who were unable to confirm earlier reports of rapid periodic changes, and by Božić et al. (1999) who found two episodes of the light brightening and identified OT Gem as the object with a positive correlation. This agrees with the inclination estimate of about 20° to 30° by Sigut & Ghafourian (2023). Balona & Ozuyar (2021) analyzed small-amplitude rapid light changes for a number of Be stars using TESS observations. For OT Gem they found a period of 0^d4606 with a non-sinusoidal light curve. Labadie-Bartz et al. (2022) – investigating two sectors of TESS photometry – noted the light brightenings and changes on longer time scales but reported also two groups of short periodic changes near 0^d249 and 0^d472.

Complete Hvar photometry is in Fig. 20 and shows frequent episodes of light brightenings, similar to V442 And.






Fig. 20: The *UB* time variations of OT Gem.

β CMi = 58715 Dulaney et al. (2017) announced that the object is a single-line spectroscopic binary with a 170^d4 period and a semiamplitude of RV curve of 2.25 km s⁻¹. However, Harmanec et al. (2019) – measuring RVs on many high-dispersion spectra – were unable to confirm their result and


Fig. 21: The *UBV* time variations of β CMi.

Fig. 22: The *UBV* time variations of κ Dra.

also Klement et al. (2024) did not detect any secondary 670
their interferometry. Božić et al. (1982) were unable to find evi-
dence of any rapid light variations, but Saio et al. (2007) de-
tected very low light variations in the MOST satellite photom-
etry, which they interpreted as g-mode nonradial pulsations.
Using wavelet analysis, Harmanec et al. (2019) reanalysed the
MOST photometry and found only one stable period of 0^d6169,
which they tentatively identified with the star rotational period.
Balona & Ozuyar (2021) found a period of 0^d6207 with a non-
sinusoidal light curve from TESS photometry.

A rather limited set of Hvar photometry is shown in Fig. 21.
There is little evidence of any changes on longer time scales. 680

BR CMi = HD 61273 A detailed spectroscopic and photo-
metric study, based largely on Hvar photometry, have been pub-
lished by Harmanec et al. (2015). BR CMi was found to be a
B9.5e + G8III semidetached binary in the late phase of the mass
transfer, having ellipsoidal light variations and a slowly increas-
ing orbital period of 12^d919. No secular or cyclic light changes
were found.

UX Mon = HD 65607 This binary was a subject of a very
detailed spectroscopic and photometric study, based also on
Hvar photometry (Sudar et al. 2011). It was found to be a rare
binary system observed at the early stage of mass exchange, be-
fore the mass ratio reversal. Its 5^d9044 orbital period was found
to be slowly decreasing with time at a rate of 0.26 s per year. In
their discussion of W Ser class of binaries and double periodic
variables Mennickent et al. (2016) claimed that the period of UX
Mon is constant. 660

HD 81357 = MWC 859 This object was found to be a
semidetached binary in the late stage of the mass exchange
and an ellipsoidal variable. Its detailed study was published by

Koubský et al. (2019) and profits from Hvar photometry. Neither
secular nor cyclic light changes were found. 690

κ Dra = 5 Dra = HD 109387 This bright Be star ob-
served in an unusually high northern declination, was found to
be a single-line spectroscopic binary with a 61^d55 orbital period
and a small RV amplitude by Juza et al. (1991). This was con-
firmed by subsequent orbital solutions by Saad et al. (2005) and
Saad et al. (2021). However, all attempts to detect the spectrum
of the secondary failed. The secondary was finally detected from
near-IR interferometry by Klement et al. (2022a). They found
a mass ratio of 0.117 ± 0.009 , in accord with the range esti-
mated by Juza et al. (1991). Juza et al. (1994) investigated spec-
tral, polarimetric and photometric observations from the past 100
years, including their own, and argued that all observables var-
ied with a period of 8406 days. Saad et al. (2004) concluded that
this variability (which they estimated to 8044 ± 167 days) is
probably cyclic, not strictly periodic. This is definitively con-
firmed by the extended series of Hvar photometry, shown in
Fig. 22, which shows a long systematic brightness decrease
since about JD 2451000 up to the present time. This nicely
correlates with the disappearance of the Balmer emission docu-
mented over a similar time interval by Klement et al. (2022a).
Balona & Ozuyar (2021) reported rotational modulation with a
period of 1^d134 from their analysis of TESS photometry. 690

V839 Her = 4 Her = HD 142926 This Be star belongs
to a few Be stars systematically monitored at Hvar since
1972. It was identified as a single-line spectroscopic binary
with a 46^d2 period (Harmanec et al. 1973b; Heard et al. 1975;
Harmanec et al. 1976). Božić et al. (2013a) reported very mild
light variations with the orbital period. The time plot of all
Hvar and transformed H_p photometric observations is in Fig. 23.
So far, the only large light decrease, associated with the forma-
tion of a new shell phase, was discovered by Percy et al. 700




Fig. 23: The UBV time variations of V839 Her. The blue dots denote observations published by Percy et al. (1997).

(1997) and is also recorded in the H_p photometry. Since this is a very interesting case, we made an exception and included also the observations by Percy et al. (1997) into our time plot (shown by blue circles). This identifies V839 Her as an object with the inverse correlation between the light and emission-line strength (Koubský et al. 1997). This is corroborated by the study of Sigut & Ghafourian (2023). These authors compared the inclination angles of a number of Be stars based alternatively on the effects of gravitational darkening, and on the $H\alpha$ emission line profile modelling. They found that V839 Her is seen almost equator-on. In photometry, no secular variations outside the shell phase have been detected in either the V magnitude or $B-V$ index, but are obvious in the $U-B$ index.

δ Sco = 7 Sco = HD 143275 This early B star was for a long time considered to be a non-variable object and was even recommended as a $uvby\beta$ standard star. Carefully analyzing earlier records and using the MAPPIT optical interferometer, Bedding (1993) discovered that the object is a spectroscopic binary with a highly eccentric orbit and a 10.5 yr orbital period and estimated that the secondary is for $1^m5\pm0^m3$ fainter than the primary. This was then confirmed by several other studies. Tycner et al. (2011) published the revised orbit with a period of 10.817 yr, eccentricity of 0.9380 and orbital inclination of 32.9° . Cote & van Kerkwijk (1993) discovered the presence of $H\alpha$ emission and since then the object became a topic of many detailed studies. Large brightness and colour variations were observed but no clear proof that the occurrence of the $H\alpha$ emission was related to the periastron passages was found. In fact a strong $H\alpha$ emission persist for a long time. Disk evolution during three periastron passages was discussed by Rast et al. (2024), who also show the light and colour curve. It is rather surprising that all attempts to find the spectral lines of the secondary failed.






Fig. 24: The UBV time variations of δ Sco.

Hvar observations are presented in Fig. 24 and complement existing observations from other stations, for instance the $B-V$ photometry by Jones et al. (2013). Note that the object was inevitably observed at large air masses from Hvar.

ζ Oph = 13 Oph = HD 149757 This star was used to be a photometric standard star in the past. However, several $H\alpha$ emission episodes were recorded in its spectra (see Ebbets 1981, and references therein). This very rapid rotator is also the first Be star for which rapid line profile variations were discovered by Walker et al. (1979). Harmanec (1989) and Harmanec (1999) interpreted the line-profile and small light changes as evidence of corotating structures revolving with a period of 0^d64 but many investigators understand these changes as evidence of non-radial pulsations. This was also the conclusion of a study based on the MOST satellite photometry (Walker et al. 2005). Hvar photometry shows only small changes and no secular variations but we warn that the star was always inevitably observed at large air masses at Hvar.

720

V744 Her = 88 Her = HD 162732 This Be star has also been systematically monitored at Hvar since 1972. It was found to be a single-line spectroscopic binary with an 86^d7 period (Harmanec et al. 1972b,a, 1974; Doazan et al. 1982b,a). Figure 25 shows the light and colour changes of the object over 50 years of Hvar observations. It is seen that the star underwent three episodes of secular light decreases, accompanied by the emission and metallic shell episodes, the first one being discussed in detail by Harmanec et al. (1978). It is a very good example of an inverse correlation between the brightness and emission-line strength (see Fig. 53). It is also seen that there is a slight secular brightness decrease at the light maxima over the whole interval of Hvar observations on the fourth time scale. There are almost no secular colour changes in $B-V$ but very


Fig. 25: The *UBV* time variations of V744 Her.

Fig. 26: The *UBV* time variations of V974 Her.


pronounced ones in $U-B$. The secondary star is probably a hot and small object but it has never been directly observed (see, e.g. Wang et al. 2018).

V974 Her = HD 164447 This is a relatively little studied B8e star. Hipparcos H_p photometry (Perryman & ESA 1997) shows a slow decrease in brightness over the whole interval of observations. As seen in the BeSS database (Neiner et al. 2011), the star lost its Balmer emission in 2020. Hvar photometry shown in Fig. 26 indicates a brightening characteristic for a positive correlation, which is rather rare for late type B stars.

σ Her = HD 166014 This B9.5e star was proposed by Perry et al. (1987) as a bright *uvbyH β* photometric standard. All the $H\alpha$ spectra in the BeSS database (Neiner et al. 2011), covering the time interval from 2011 to 2024, show weak double $H\alpha$ emission not reaching the continuum level. A limited set of Hvar photometry shown in Fig. 27 is indicative of small rapid changes and no secular variations.

NW Ser = HD 168797 The Hvar photometry of this B2e star was analysed together with photometry secured by John R. Percy in Toronto to document its rapid and possibly periodic light changes on time scales from 0^d.4 to 5^d.5 days (Percy et al. 1999). No more recent Hvar observations were secured since the publication of that study. Fig. 28 shows the time plot of all available Hvar and transformed H_p observations.

CX Dra = HD 174237 This object is now known as a B2e+F5III semi-detached binary with a 6^d.696 period seen under an intermediate inclination of the orbit – see the RV studies by Koubský (1976, 1978); Horn et al. (1992); Richards et al. (2000), and references therein. The light and colour variations show cyclic changes correlated with the variations of the strength


Fig. 27: The *UBV* time variations of σ Her.

of the $H\alpha$ emission on a time scale of several hundred days, showing a positive correlation (Koubský et al. 1980, 1998). Besides that, there are low-amplitude phase-locked light changes with rather unusual properties – see Fig. 3 in Koubský et al. (1998). The lower envelope of these changes is reminiscent of




Fig. 28: The *UB* time variations of NW Ser.




Fig. 29: The *UB* time variations of CX Dra.

ellipticity and reflection in the undisturbed state, but frequent brightenings are seen. We show the complete light and colour changes recorded at Hvar in Fig. 29.

β Lyr = 10 Lyr = HD 174638 This is one of two first discovered Be stars and a famous 12^d.93 eclipsing and strongly






Fig. 30: The orbital light curve of β Lyr.

interacting binary with a complicated structure of circumstellar matter comprising from a disk, scattering halo and bipolar jets and with a rapidly increasing orbital period (19 s per year). The history of its investigation was summarized by Harmanec (2002). Hvar observations (shown in a phase plot Fig. 30) have been secured in support of several large studies, which also provide summaries of more recent studies: Harmanec et al. (1996); Ak et al. (2007); Mourard et al. (2018); Brož et al. (2021). For a long time known cycle-to-cycle variations inside the binary eclipses are seen there. There is, however, no secular trend in either the brightness in *V* passband or in the colour indices.

V923 Aql = HD 183656 This Be star was found to be the primary component of a 214^d.7 binary by Koubský et al. (1989). Its very detailed study, based on all available spectral and photometric observations over many decades was published by Wolf et al. (2021). They documented its long-term, orbital and rapid variations. The object is a typical example of the inverse correlation between the brightness and emission-line strength. While it exhibits large-amplitude cyclic RV and *V/R* variations


Fig. 31: The *UBV* time variations of V923 Aql.

Fig. 32: The *UBV* time variations of V1294 Aql.

with cycle lengths between about 1800, and 3000 days in RV, V/R , V magnitude, and $U-B$ index, its $B-V$ index remains secularly stable and close to zero. Rapid changes, if periodic, may follow a 0^d8442 period. In Fig. 31 we show the time plot of Hvar photometry, extended a bit for more recent observations. No evidence of changes on the fourth, longest time scale is seen.

V1294 Aql = HD 184279 This one of the Be stars with the largest recorded range of light variations. Its amazing and complicated spectral, light and colour variations were recently described in detail in the study by Harmanec et al. (2022), who discovered that the object is also a spectroscopic binary with a 192^d9 period. The readers are referred to this study. Here we only show the extended series of photometric observations from Hvar in Figs. 32. It is seen that the brightness of the object is still rising and is maximal over the whole recorded history.

V1507 Cyg = HD 187399 This object was known to have $H\alpha$ emission since 1921 and was found to be a peculiar binary with an eccentric orbit and a 27^d97 orbital period (Merrill 1949). Hutchings & Redman (1973) interpreted the object as a binary in the phase before the onset of a large scale mass exchange and discussed also a very peculiar orbital light curve, which was later published in detail by Hill et al. (1976). The peculiar shape of the light curve was confirmed by our Hvar observations (Pavlovski et al. 1979). Hutchings & Redman (1973) interpreted the light-curve minima as evidence of shallow eclipses. Davidge (2023) published a very detailed study of the binary and also summarized all other published studies. Measuring weak emission wings of the Si II 6347 and 6371 Å lines, similarly as several investigators did for β Lyr, he obtained a RV curve of the secondary and estimated the component masses. He argued that the brighter primary, whose spectrum is closely reminiscent of the spectra of bright supergiants such as β Ori, is now

the less massive star of the system, which lost its mass toward the secondary during the earlier rapid phase of the mass transfer. He correctly pointed out that the orbital period of 27^d9705 (Hutchings & Redman 1973) has remained constant within the limits of the accuracy of its determination since Merrill (1949) study, which implies that the system must now be in the stage of terminal, slow mass transfer. He also speculated about the possibility that the unusually eccentric ($e=0.388$) orbit could be the consequence of perturbation by a third star in the system or fly-by. Koenigsberger & Estrella-Trujillo (2024) investigated the role of the tidal energy dissipation for stars undergoing light brightenings during periastron passage and modelled such a effect.

The orbital light curve for the combined Hvar and Hipparcos H_p observations transformed to Johnson V is shown in Fig. 33. No secular light changes were found.

V1624 Cyg = 28 Cyg = HD 191610 This Be star is well-known for its pronounced rapid photometric variations (Percy & Lane 1977; Mills et al. 1979; Pavlovski et al. 1997, for instance). Baade et al. (2018) analyzed long strings of photometry from the BRITE and SMEI satellites in relation to the long-term variations of the strength of the $H\alpha$ emission. They identified a number of rapid periods ranging from 0^d318 to 0^d759, some of them being combinations of others, and also a period of 19^d634 with the largest amplitude, seen in both, SMEI and BRITE data. They advocated the hypothesis that the coupled non-radial pulsation are connected to the mass transfer from the stellar photosphere to the Be star disk. Wang et al. (2018) tried to detect new hot subdwarf companions to Be stars in the IUE spectra. For V1624 Cyg they tabulated RVs for 25 spectra. Wang et al. (2021) investigated far-UV spectra from the Imaging Spectrograph of the Hubble Space telescope (HST/STIS) and were unable to find any lines of the secondary. Klement et al. (2022b) observed the object interferometrically




Fig. 33: The 27^d9705 orbital light curve of V1507 Cyg based on Hvar (black) and transformed Hipparcos H_p (red) observations.

with the CHARA array and tentatively suggested an orbit with a 246 d period. Later, Klement et al. (2024) concluded from extended interferometry covering about one half of the orbital period that the true binary has a circular orbit with a 359^d260 period and orbital inclination of 119° (i.e. 61°). Using RVs from Wang et al. (2018) and Wang et al. (2021) papers and the estimated distance to the system, they estimated the masses of both stars. Sigut & Ghafourian (2023) estimated the inclination of the Be disk from H α emission profile fitting as $40^\circ \pm 5^\circ$ and $69^\circ \pm 17^\circ$ from the gravity darkening so it is possible that the disk lies on the orbital plane of the binary.

Hvar photometry is shown in Fig. 34. It is seen that the variations are dominated by rapid changes. There are obviously also changes on longer time scales but our limited set of observations does not allow to do any firm conclusions about their character.

20 Vul = HD 192044 This is a little studied B8 star. Copeland & Heard (1963) reported broad nebulous H α and He I




Fig. 34: The *UBV* time variations of V1624 Cyg.





Fig. 35: The *UBV* time variations of 20 Vul.

lines and double H β and H γ emission lines with variable intensity and V/R changes for their photographic spectra from the years 1938 to 1961 but no details were given. We are not aware of any systematic study devoted this object. Available Hvar photometry is shown in Fig. 35.


Fig. 36: The *UBV* time variations of QR Vul.

QR Vul = HR 192685 This object is the brighter component of a visual binary ADS 13589, the component B located only $0''.7$ from it and being for $2^m.7$ fainter. Balona & Ozuyar (2021) analyzed the TESS observations and suggested that the light variations can be phased with a $0^d.7918$ period with a non-sinusoidal shape, which they identified with the period of rotation.

Limited Hvar photometry shown in Fig. 36 is indicative of rapid changes and occasional light brightenings.



P Cyg = 34 Cyg = HD 93237

Brightness variations of this famous luminous blue variable have been monitored for 5 years with the BRITE satellites and the results are presented by Elliott et al. (2022). These authors were unable to find any consistent periodicities and concluded that the variations are stochastic in their nature.

Our limited Hvar observations are displayed in Fig. 37. They can complement the existing observations. We note that the variations in the $U-B$ index are larger than those in $B-V$.



25 Vul = HD 193911 Balona & Ozuyar (2021) concluded from the TESS photometry that the small light changes can be reconciled with a $1^d.890$ period with a non-sinusoidal light curve. They identified it with the star rotational period. Limited Hvar photometry shown in Fig. 38 is indicative of small-amplitude rapid changes, with possibly some secular evolution in the $U-B$ index..

V2119 Cyg = HD 194335 Klement et al. (2024) obtained an interferometric circular orbit with a period of $63^d.1475$ and summarised the previous studies of this be binary with a hot compact companion. Regrettably, no optical RV curve is available.


Fig. 37: The *UBV* time variations of P Cyg.

Fig. 38: The *UBV* time variations of 25 Vul.

Hvar photometry is shown in Fig. 39. A plot of combined Hvar V magnitude and transformed H_p magnitude versus the orbital phase does not show a convincing light curve.


V1661 Cyg = 55 Cyg = HD 198478 This star is classified as a B3-4Ia supergiant. It was a subject of almost 500

Fig. 39: The *UBV* time variations of V2119 Cyg.Fig. 40: The *UBV* time variations of V1661 Cyg.



studies so we only refer readers to the paper of Kraus et al. (2015), where the studies of spectral and photometric variability of V1661 Cyg are summarised. These authors tried to study an interplay between the mass loss and pulsations and decomposed the observed variations in several observables into a number of periods ranging from 2.7 hours to 22^d5.

Our limited set of Hvar observations is shown in Fig. 40. It shows some variability in *V* and *B*−*V* colour but, notably, no variability in the *U*−*B* index.

V832 Cyg = 59 Cyg = HD 200120 This object is the brightest component A of the multiple visual system ADS 14526, with a rather distant fainter components B, C, D, and E. Already Harmanec (1982) speculated that – at that time already known – spectral variations and appearance of shell phases of V832 Cyg could be related to the presence of as yet unknown close visual companion. McAlister et al. (1984) indeed discovered such a companion Ab and its orbital variations has been monitored since then. Mason published a preliminary orbit of the Aa-Ab pair with an orbital period of 161.5 yr and eccentricity of 0.261 in Information Circular 175 of the IAU Comm. 26, but no details were given. Tarasov & Tuominen (1987) discovered radial-velocity (RV) and *V*/*R* variations of the H α emission with a 29^d14 period, which they interpreted as the evidence that the object is a spectroscopic binary. Harmanec et al. (2002) carried out a detailed study of this Be stars, based on spectral and photometric observations from several observatories. They indeed confirmed that V832 Cyg is a spectroscopic binary with a 28^d1971 period and argued that the true orbital RV amplitude is best defined by the RVs measured on the steep emission wings of the H α line profile. This was nicely confirmed by the study of Peters et al. (2013), who derived the RV curves of both binary components from the far-UV spectra and proved that the secondary is a hot O type subdwarf. Harmanec et al. (2002) also documented the

Fig. 41: The *UBV* time variations of V832 Cyg.

long-term light and colour changes with a positive correlation between the light and emission-line strength combined with changes probably attributable to one-arm disk oscillation. Prewhitening photometry for these long-term changes, they found mild sinusoidal light changes with the orbital period.


Fig. 42: The *UBV* time variations of V1931 Cyg.

Fig. 43: The *UBV* time variations of V382 Cep.

The long-term brightness and colour variations of V832 Cyg extended to the present time are shown in Fig. 41 and the accompanying behaviour in the colour-colour diagram is in Fig. 52. One can see a steady rise of the brightness over the whole time interval covered by the *UBV* observations, attributable to the variability on the fourth time scale.

V1931 Cyg = 60 Cyg = HD 200310 This object is the brighter component of a visual binary WDS J21012+4609. Koubský et al. (2000) discovered that the object is also a spectroscopic binary with a 146^d6 period and analysing photometry, including Hvar observations, they reported large cyclic changes and several possible rapid periods near 1 d. For comparison, Balona & Ozuyar (2021) reported a rotational period of 0^d6109 with a double-wave light curve from TESS photometry. The binary was resolved interferometrically and the combined spectroscopic and astrometric solution gave an improved period of 147^d617, an eccentric orbit with $e = 0.20$, and the orbital inclination of 84°6 (Klement et al. 2024). Hvar photometry plotted in Fig. 42 shows the pronounced cyclic light and colour changes on a time scale of years. Surprisingly, it shows a clear positive correlation in the $U-B$ vs. $B-V$ diagram. Is the circumstellar disk tilted with respect to the binary orbit?

1040

V382 Cep = 6 Cep = HD 203467 Krajčeva & Koubský (1983) analyzed two sets of RVs from photographic spectra and suggested that the object could be a close spectroscopic binary with periods of either 2^d4389 or 0^d7085. Balona & Ozuyar (2021) found a photometric period of 1^d658, which they identified with the star rotational period. Limited Hvar observations are shown in Fig. 43. They show some secular changes but little can be said about their character.

8 Lac A = HD 214168 = HIP 111546 8 Lac is a multi-component system WDS J22359+3938 composed of components Aa,

Ab, B, C, D, E, F, G, H, I, and J. Component B is also an early B star with $V = 6^m48$ according to a few Hvar observations, in agreement with the HD catalog. We note that the SIMBAD bibliography gives incorrectly $V = 5^m67$ for 8 Lac B. Regrettably, there are no Hipparcos H_p observations with error flags 0 or 1 in the Hipparcos catalogue for either 8 Lac A or 8 Lac B. Balona & Ozuyar (2021) classify the star as β Cep and rotational variable from TESS photometry and give a rotation period of 0^d4277. Klement et al. (2024) were unable to resolve the binary in a consistent way from their interferometric observations.

Hvar photometry is shown in Fig. 44. It indicates mild rapid and secular cyclic changes but little can be concluded about their character.




1000

HR 8682 = HD 216057 This B8 star has no variable-star name and the BeSS spectra since 2001 to 2024 show no or only marginal $H\alpha$ emission (Neiner et al. 2011). A limited set of Hvar photometry is in Fig. 45. No clear evidence of variability was found.

V360 Lac = HD 216200 This object was found to be a semi-detached Be+F binary with a critically rotating secondary and an orbital period of 10^d085 (Hill et al. 1997; Linnell et al. 2006). There are also mild cyclic brightness and colour changes but no slow changes on the fourth time scale. Since no new Hvar photometry was obtained since the publication of Linnell et al. (2006), we refer readers to the two cited papers, where also other studies of this object are summarised.

EW Lac = HD 217050 This is a very active Be star, worth of a detailed and complex study. It has been studied for about 100 years. Its Balmer emission was variable until 1926. Then a strong emission with sharp shell lines visible over the whole Balmer series has developed and remained stable until 1977, when the

1050


Fig. 44: The *UBV* time variations of 8 Lac A.

Fig. 46: The *UBV* time variations of EW Lac.

Fig. 45: The *UBV* time variations of HR 8682.



detected in previous studies, but with a light curve variable with time. The spectra from the BeSS database (Neiner et al. 2011) show that the emission gradually weakened and is virtually missing since 2022.

Complete Hvar photometry shown in Fig. 46 indicates that the star is an example of variability on the fourth time scale, with secular brightness increase. It is also seen that the amplitude of rapid variations decreased as the star was gradually losing the Balmer emission.

o And = HD 217675 = HR 8762 This is a remarkable Be star, for which rapid photometric variability was discovered already during World War I, based on photoelectric observations published by Guthnick & Prager (1918). Later, Guthnick (1941) reported a 1^d5765 period of light variations and the star was long considered to be a close contact binary (see Harmanec 1983, 1070 for the whole early history of the search for the true period of its changes). Later, it was found that *o* And is a multiple system, containing a double-lined spectroscopic binary with a 33^d085 period, moving in a wide orbit with the Be star, which dominates the optical spectrum (Hill et al. 1988; Harmanec et al. 1987a). The object exhibits rapid light variations with a period of $1^d571272$ and a variable shape of the light curve (Harmanec 1984a; Harmanec et al. 1987b; Stagg et al. 1988) and superimposed cyclic variation with a regular cycle length over 2500 d (8.5 yr). Harmanec (1984a) suggested it could be related to a visual binary discovered by Wilson (1950) and causing recurrent shell phases. Hvar photometry shown in Fig. 47 shows regular changes with a possible period of 2525 d.

star became active again (see, e.g. Harmanec et al. 1979). These authors also showed that the star is a classical example of the positive correlation between the brightness and emission-line strength. Stagg et al. (1988) summarised the history of its photometric studies and suggested a period of 0^d72 , close to timescales

KX And = HD 218393 A large spectroscopic and photometric study, based also on Hvar photometry, was published by Štefl et al. (1990). These authors found periodic RV, light and colour variations with a 38^d919 period and suggested tentatively that the object could be an interacting binary. Floquet et al.


Fig. 47: The *UBV* time variations of *o* And.

Fig. 48: The *UBV* time variations of KX And.

(1995) and Tarasov et al. (1998) obtained a clear circular-orbit period based on sharp lines of G8II spectrum with a semiamplitude of 86 km s^{-1} , proving thus definitively the binary nature of the object.

Hvar photometry is in Fig. 48. No secular changes are seen. The orbital light curves are in Fig. 49.

KY And = HD 218674 Stagg et al. (1988) published the results of an international observing spectral and photometric campaign on five Be stars including KY And. Hvar photometry represented a significant contribution. While rapid light variations were undoubtedly found for KY And, determination of a consistent period or a combination of several periods was not convincing. The most promising period of 0^d753 was detectable only in a part of the data.

Complete Hvar photometry is in Fig. 50. It is obvious that the rapid changes dominate, no secular trends are seen.

LQ And = HD 224559 This is one of the first Be stars for which the presence of small-amplitude periodic light variations was found (Percy 1983). A very detailed period analysis of numerous photometric observations, including numerous observations from Hvar by Harmanec et al. (1991) led to the conclusion that that light variations follow a 0^d61904 period with a double-wave light curve, which was tentatively identified with the star rotation period. Matthews et al. (1991) analyzed a series of electronic spectra and reported that the object could be a single-line spectroscopic binary with a 7^d413 orbital period. The RV residuals from the orbital solution as well as line intensities and equivalent widths were found to vary with the photometric 0^d61904 period. It is necessary to add that the binary period was not confirmed in any of the follow-up studies. The unpublished $H\alpha$ spectra and spectra from the BeSS database indicate that during the past 20 years the star had a strong emission reaching up to 4 times the continuum level.

Hvar photometry is shown in Fig. 51.

4. Discussion

Our systematic observational effort and monitoring of a representative sample of bright Be stars from the Northern Hemisphere (down to declinations of about -20°) helped to realization that virtually all Be stars are light and colour variables and to the recognition of several of their characteristic timescales.

With the exception of several dedicated observational campaigns, our photometry is more suited to the detection of changes on long time scales.

The cases of objects like β Lyr or V1507 Cyg show how the duplicity or multiplicity of individual emission-line objects can lead to a large variety of observational effects. In particular, the Roche-lobe filling brighter but usually less massive secondaries in the binaries with large-scale mass transfer can have spectra, which closely mimic the supergiant or bright giant spectral classification but have masses significantly lower than $10 M_\odot$.

It is also notable that the vast majority of known binaries with the Be primary has circular orbits. This seems to indicate that these systems underwent large scale mass transfer sometimes in their evolution, which helped to circularize their orbits. For V1507 Cyg, which has an eccentric orbit, Davidge (2023) argued that the eccentricity was caused by a third body in the system. This could also be the case of BU Tau, a member of the Pleiades cluster, where the encounter with another body seems quite probable. Also another object with an eccentric orbit, V1931 Cyg, is a member of a relatively close visual system.

Our sample indicates that the least variable Be stars are found among objects with spectral types later than B5. Shokry et al. (2018) – who studied a different sample of Be stars – arrived at a similar conclusion. However, this is not a general rule. Objects like V839 Her, V744 Her or BU Tau actually show large secular




Fig. 49: The 38^d919 orbital light curve of KX And based on Hvar (black) and transformed Hipparcos H_p (red) observations.

changes. Could that be related to the fact that all of them are known binaries?

We note that in some cases of secular changes, the amplitude of variations in the $B-V$ index is larger than that in $U-B$, while in other objects it is just opposite.

The existence of variations on the four time scale implies that it is not easy to estimate the true stellar radius from the dereddened V magnitude, bolometric correction and measured parallax since even the observed V magnitude outside the emission-line episodes varies with time and probably refers to the remaining stellar pseudophotosphere.

We wish to issue a warning that it can be somewhat dangerous to built models of Be star disks based only on satellite photometries with their very broad passbands only. Cases of stars like V744 Her, V1294 Aql or BU Tau show that the variations in the yellow, blue, and ultraviolet parts of the stellar spectrum can be quite different and these differences are inevitably affecting




Fig. 50: The UBV time variations of KY And.




Fig. 51: The UBV time variations of LQ And.

the variability patterns recorded with the very broad passbands of the MOST, TESS, and largely also BRITE satellites.

In Figures 52 and 53 we show examples of the positive and inverse correlations between the brightness and Balmer-line emission strength as defined in Sect. 3. We note that besides the




Fig. 52: The $U-B$ versus $B-V$ diagram for several Be stars with a positive correlation between the brightness and emission-line strength.




Fig. 53: The $U-B$ versus $B-V$ diagram for several Be stars with an inverse correlation between the brightness and emission-line strength.

basic type, the inverse correlation is also observed on the time scale of binary orbital motion for binaries observed not far from the equator-on configuration. This is obviously due to varying projection effects of circumstellar matter as these binaries revolve around each other. This is the case of KX And, UX Mon or RX Cas in Fig. 53.

In passing, we would like to admit openly that we have deliberately postponed the publication of the complete Hvar data archives for a late date. The reason is that we still plan several dedicated studies of objects studied more systematically at Hvar. We are, however, open to sharing the data with colleagues seriously interested in collaboration with us on such studies.

Acknowledgements. H.B. acknowledges financial support from Croatian Science Foundation under the project 6212 “Solar and Stellar Variability”. P.H., M.W.,

M.B., A.O., J.J., and K.V. were supported by the Czech Science Foundation grant GA19-01995S. Our thanks go to our colleagues and students A. Arena, J. Arsenijević, R. Brajša, R. Brož, P. Chadima, D. Čikotić, E. Frlež, J. Grygar, K. Hoňková, Z. Ivanović, Ž. Ivezić, J. Janík, J. Juryšek, B. Jovanović, G. Lazin, M. Malarić, M. Mulić, M. Muminović, M. Netopil, D. Ondřích, I. Piantschitsch, D. Plačko-Vršnak, P. Polechová, Ž. Ružić, M. Stupar, I. Vince, B. Vršnak, and M. Zejda who obtained some Hvar observations reported here and are not included among the co-authors. Our special thanks go to J. Vacek for his design, maintenance and regular servicing of the Hvar photoelectric photometer. J.R. Percy kindly put the archive of his *UBV* observations of bright Be stars at our disposal. The following internet-based resources were consulted: the SIMBAD database and the VizieR service operated at CDS, Strasbourg, France; and the NASA’s Astrophysics Data System Bibliographic Services.

References

- Abt, H. A., Barnes, R. C., Biggs, E. S., & Osmer, P. S. 1965, *ApJ*, 142, 1604
Ak, H., Chadima, P., Harmanec, P., et al. 2007, *A&A*, 463, 233
Alvarez, M. & Schuster, W. J. 1981, *Rev. Mexicana Astron. Astrofis.*, 6, 163
Andersen, J., Pavlovski, K., & Pirola, V. 1989, *A&A*, 215, 272
Baade, D., Pigulski, A., Rivinius, T., et al. 2018, *A&A*, 610, A70
Ballereau, D. & Chauville, J. 1989, *A&A*, 214, 285
Balona, L. A., Aerts, C., Božić, H., et al. 2001, *MNRAS*, 327, 1288
Balona, L. A. & Engelbrecht, C. A. 1986, *MNRAS*, 219, 131
Balona, L. A. & Ozuyar, D. 2021, *ApJ*, 921, 5
Barnsley, R. M. & Steele, I. A. 2013, *A&A*, 556, A81
Bedding, T. R. 1993, *AJ*, 106, 768
Bemporad, A. 1904, *Mitteil. Grossherzog. Sternwarte Heidelberg*, 4, 1
Bergin, E. A., Burns, J. F., Guinan, E. F., & McCook, G. P. 1989, *Information Bulletin on Variable Stars*, 3358, 1
Bolton, C. T. 1982, in *Be Stars*, ed. M. Jaschek & H. G. Groth, Vol. 98, 181–184
Borre, C. C., Baade, D., Pigulski, A., et al. 2020, *A&A*, 635, A140
Božić, H., Ruždjak, D., & Sudar, D. 1999, *A&A*, 350, 566
Božić, H. & Harmanec, P. 1998, *A&A*, 330, 222
Božić, H., Harmanec, P., Horn, J., et al. 1995, *A&A*, 304, 235
Božić, H., Harmanec, P., & Koubský, P. 2013a, *Central European Astrophysical Bulletin*, 37, 9
Božić, H., Harmanec, P., & Koubský, P. 2013b, *Central European Astrophysical Bulletin*, 37, 9
Božić, H., Harmanec, P., Yang, S., et al. 2004, *A&A*, 416, 669
Božić, H., Muminović, M., Pavlovski, K., et al. 1982, *Information Bulletin on Variable Stars*, 2123, 1
Božić, H. & Pavlovski, K. 1988, *Hvar Observatory Bulletin*, 12, 15
Breger, M. 1972, *ApJ*, 176, 367
Brož, M., Mourard, D., Budaj, J., et al. 2021, *A&A*, 645, A51
Burki, G., Rufener, F., Burnet, M., et al. 1995, *A&AS*, 112, 383
Carciofi, A. C., Okazaki, A. T., Le Bouquin, J. B., et al. 2009, *A&A*, 504, 915
Copeland, J. A. & Heard, J. F. 1963, *Publications of the David Dunlap Observatory*, 2, 317
Cote, J. & van Kerkwijk, M. H. 1993, *A&A*, 274, 870
Cousins, A. W. J. & Jones, D. H. P. 1976, *MNRAS*, 81, 1
Davidge, T. J. 2023, *AJ*, 166, 188
Delplace, A. M. 1970, *A&A*, 7, 68
Doazan, V., Harmanec, P., Koubský, P., Krpata, J., & Žďárský, F. 1982a, *A&A*, 115, 138
Doazan, V., Harmanec, P., Koubský, P., Krpata, J., & Žďárský, F. 1982b, *A&AS*, 50, 481
Dorren, J. D., Fu, H. H., & Wu, H. H. 1984, *Information Bulletin on Variable Stars*, 2521, 1
Dulaney, N. A., Richardson, N. D., Gerhart, C. J., et al. 2017, *ApJ*, 836, 112
Ebbets, D. 1981, *PASP*, 93, 119
Elliott, A., Richardson, N. D., Pablo, H., et al. 2022, *MNRAS*, 509, 4246
Floquet, M., Hubert, A. M., Hubert, H., Ballereau, D., & Chauville, J. 1995, *A&A*, 294, 227
Gaposchkin, S. 1944, *ApJ*, 100, 230
Ghoreyshi, M. R., Carciofi, A. C., Jones, C. E., et al. 2021, *ApJ*, 909, 149
Ghoreyshi, M. R., Jones, C. E., & Granada, A. 2023, *MNRAS*, 518, 30
Gies, D. R., Bagnuolo, William G., J., Ferrara, E. C., et al. 1998, *ApJ*, 493, 440
Gies, D. R., McKibben, W. P., Kelton, P. W., Opal, C. B., & Sawyer, S. 1990, *AJ*, 100, 1601
Gies, D. R., Wang, L., & Klement, R. 2023, *ApJ*, 942, L6
Gies, D. R., Willis, C. Y., Penny, L. R., & McDavid, D. 1993, *PASP*, 105, 281
Grundstrom, E. D. 2007, *PhD thesis*, Georgia State University
Guinan, E. F. & Hayes, D. P. 1984, *ApJ*, 287, L39
Guthnick, P. 1941, *Vierteljahrsschrift Astron. Gesell.*, 76
Guthnick, P. & Prager, R. 1918, *Veröff. Berlin Babelsberg*, 2, 113
Harmanec, P. 1980, *Be Star Newsletter*, 1, 2

- Harmanec, P. 1982, in *Be Stars*, ed. M. Jасhek & H. G. Groth, Vol. 98, 279–297
- Harmanec, P. 1983, *Hvar Observatory Bulletin*, 7, 55
- Harmanec, P. 1984a, *Information Bulletin on Variable Stars*, 2506, 1
- Harmanec, P. 1984b, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 35, 193
- 1270 Harmanec, P. 1989, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 40, 201
- Harmanec, P. 1994, in *NATO Advanced Study Institute (ASI) Series C*, Vol. 436, *The Impact of Long-Term Monitoring on Variable Star Research: Astrophysics*, ed. C. Sterken & M. de Groot, 55
- Harmanec, P. 1998a, *A&A*, 335, 173
- Harmanec, P. 1998b, *A&A*, 334, 558
- Harmanec, P. 1999, *A&A*, 341, 867
- Harmanec, P. 2000, in *Astronomical Society of the Pacific Conference Series*, Vol. 214, *IAU Colloq. 175: The Be Phenomenon in Early-Type Stars*, ed. M. A. Smith, H. F. Hennrichs, & J. Fabregat, 13
- 1280 Harmanec, P. 2002, *Astronomische Nachrichten*, 323, 87
- Harmanec, P. & Božić, H. 2013, *Central European Astrophysical Bulletin*, 37, 3
- Harmanec, P., Božić, H., Koubský, P., et al. 2022, *A&A*, 666, A136
- Harmanec, P., Božić, H., Percy, J. R., et al. 2002, *A&A*, 387, 580
- Harmanec, P., Grygar, J., Horn, J., et al. 1977a, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 28, 133
- Harmanec, P., Habuda, P., Štefl, S., et al. 2000, *A&A*, 364, L85
- Harmanec, P., Hill, G. M., Walker, G. A. H., Dinshaw, N., & Yang, S. 1987a, *Publications of the Astronomical Institute of the Czechoslovak Academy of Sciences*, 5, 115
- 1290 Harmanec, P., Horn, J., & Juza, K. 1994, *A&AS*, 104, 121
- Harmanec, P., Horn, J., & Koubský, P. 1980a, *Be Star Newsletter*, 2, 3
- Harmanec, P., Horn, J., & Koubský, P. 1982, in *Be Stars*, ed. M. Jасhek & H. G. Groth, Vol. 98, 269
- Harmanec, P., Horn, J., Koubský, P., et al. 1977b, *Information Bulletin on Variable Stars*, 1324, 1
- Harmanec, P., Horn, J., Koubský, P., et al. 1978, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 29, 278
- Harmanec, P., Horn, J., Koubský, P., et al. 1979, *Information Bulletin on Variable Stars*, 1555, 1
- 1300 Harmanec, P., Horn, J., Koubský, P., et al. 1980b, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 31, 144
- Harmanec, P., Horn, J., Koubský, P., et al. 1983, *Information Bulletin on Variable Stars*, 2447, 1
- Harmanec, P., Koubský, P., Horn, J., & Havelka, J. 1973a, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 24, 311
- Harmanec, P., Koubský, P., & Krpata, J. 1972a, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 23, 218
- Harmanec, P., Koubský, P., & Krpata, J. 1972b, *Astrophys. Lett.*, 11, 119
- Harmanec, P., Koubský, P., & Krpata, J. 1973b, *A&A*, 22, 337
- 1310 Harmanec, P., Koubský, P., & Krpata, J. 1974, *A&A*, 33, 117
- Harmanec, P., Koubský, P., Krpata, J., & Žďárský, F. 1976, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 27, 47
- Harmanec, P., Koubský, P., Nemravová, J. A., et al. 2015, *A&A*, 573, A107
- Harmanec, P. & Kríž, S. 1976, in *Be and Shell Stars*, ed. A. Slettebak, Vol. 70, 385
- Harmanec, P., Matthews, J. M., Božić, H., et al. 1991, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 42, 1
- Harmanec, P., Morand, F., Bonneau, D., et al. 1996, *A&A*, 312, 879
- Harmanec, P., Olah, K., Božić, H., et al. 1987b, in *IAU Colloq. 92: Physics of Be Stars*, ed. A. Slettebak & T. P. Snow, 456
- 1320 Harmanec, P., Pavlovski, K., Božić, H., et al. 1997, *Journal of Astronomical Data*, 3, 5
- Harmanec, P., Švanda, M., Korčáková, D., et al. 2019, *ApJ*, 875, 13
- Heard, J. F., Hurkens, R., Harmanec, P., Koubský, P., & Krpata, J. 1975, *A&A*, 42, 47
- Hill, G., Harmanec, P., Pavlovski, K., et al. 1997, *A&A*, 324, 965
- Hill, G., Hilditch, R. W., & Pfannenschmidt, E. L. 1976, *Publications of the Dominion Astrophysical Observatory Victoria*, 15, 1
- 1330 Hill, G. M., Walker, G. A. H., Dinshaw, N., Yang, S., & Harmanec, P. 1987b, *PASP*, 100, 243
- Hirata, R. 1995, *PASJ*, 47, 195
- Hirata, R. & Kogure, T. 1976, *PASJ*, 28, 509
- Hoffleit, D. & Jасhek, C. 1982, *The Bright Star Catalogue*. Fourth revised edition. (Containing data compiled through 1979).
- Horn, J., Harmanec, P., Koubský, P., et al. 1982, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 33, 308
- Horn, J., Hubert, A. M., Hubert, H., Koubský, P., & Bailloux, N. 1992, *A&A*, 259, L5
- Hummel, W. & Štefl, S. 2001, *A&A*, 368, 471
- 1420 Hutchings, J. B. & Redman, R. O. 1973, *MNRAS*, 163, 209
- Hutter, D. J., Tycner, C., Zavala, R. T., et al. 2021, *ApJS*, 257, 69
- Johnson, H. L. & Harris, D. L. 1954, *ApJ*, 120, 196
- Johnson, H. L., Mitchell, R. I., Iriarte, B., & Wisniewski, W. Z. 1966, *Communications of the Lunar and Planetary Laboratory*, 4, 99
- Jones, C. E., Tycner, C., & Smith, A. D. 2011, *AJ*, 141, 150
- Jones, C. E., Wiegert, P. A., Tycner, C., et al. 2013, *AJ*, 145, 142
- Juza, K., Harmanec, P., Božić, H., et al. 1994, *A&AS*, 107, 403
- Juza, K., Harmanec, P., Hill, G. M., Tarasov, A. E., & Matthews, J. M. 1991, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 42, 39
- 1350 Katahira, J.-I., Hirata, R., Ito, M., et al. 1996, *PASJ*, 48, 317
- Kazarovets, E. V., Samus, N. N., Durlevich, O. V., et al. 1999, *Information Bulletin on Variable Stars*, 4659, 1
- Klement, R., Baade, D., Rivinius, T., et al. 2022a, *ApJ*, 940, 86
- Klement, R., Rivinius, T., Gies, D. R., et al. 2024, *ApJ*, 962, 70
- Klement, R., Schaefer, G. H., Gies, D. R., et al. 2022b, *ApJ*, 926, 213
- Koenigsberger, G. & Estrella-Trujillo, D. 2024, *A&A*, 685, A145
- Koubský, P. 1976, *Information Bulletin on Variable Stars*, 1188, 1
- Koubský, P. 1978, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 29, 288
- 1360 Koubský, P., Harmanec, P., Božić, H., et al. 1998, *Hvar Observatory Bulletin*, 22, 17
- Koubský, P., Harmanec, P., Brož, M., et al. 2019, *A&A*, 629, A105
- Koubský, P., Harmanec, P., Gulliver, A. F., Ballereau, D., & Chauville, J. 1989, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 40, 31
- Koubský, P., Harmanec, P., Horn, J., et al. 1980, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 31, 75
- Koubský, P., Harmanec, P., Hubert, A. M., et al. 2000, *A&A*, 356, 913
- Koubský, P., Harmanec, P., Kubát, J., et al. 1997, *A&A*, 328, 551
- Koubský, P., Hummel, C. A., Harmanec, P., et al. 2010, *A&A*, 517, A24
- 1370 Koubský, P. & Pavlovski, K. 1982, *Hvar Observatory Bulletin*, 6, 1
- Krajčeva, Z. T. & Koubský, P. 1983, *Information Bulletin on Variable Stars*, 2313, 1
- Kraus, M., Hauke, M., Cidale, L. S., et al. 2015, *A&A*, 581, A75
- Krelowski, J., Strobel, A., Galazutdinov, G. A., Bondar, A., & Valyavin, G. 2019, *MNRAS*, 486, 112
- Kříž, S., Arsenijević, J., Grygar, J., et al. 1980, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 31, 284
- Kříž, S. & Harmanec, P. 1975, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 26, 65
- 1380 Labadie-Bartz, J., Carciofi, A. C., Henrique de Amorim, T., et al. 2022, *AJ*, 163, 226
- Linnell, A. P., Harmanec, P., Koubský, P., et al. 2006, *A&A*, 455, 1037
- Lockwood, G. W. & Thompson, D. T. 1986, *AJ*, 92, 976
- Matthews, J. M., Harmanec, P., Walker, G. A. H., Yang, S., & Wehlau, W. H. 1991, *MNRAS*, 248, 787
- Mayer, P. 1976, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 27, 308
- 1390 Mayer, P. 1977, *Hvar Observatory Bulletin*, 1, 5
- Mayer, P. 2013, *Central European Astrophysical Bulletin*, 37, 1
- McAlister, H. A., Hartkopf, W. I., Gaston, B. J., Hendry, E. M., & Fekel, F. C. 1984, *ApJS*, 54, 251
- McAlister, H. A., Hartkopf, W. I., Sowell, J. R., Dombrowski, E. G., & Franz, O. G. 1989, *AJ*, 97, 510
- McNamara, B. J. 1985, *ApJ*, 289, 213
- McNamara, B. J. 1987, *ApJ*, 312, 778
- Mennickent, R. E., Djurašević, G., Petrović, J., et al. 2022, *A&A*, 666, A51
- Mennickent, R. E., Otero, S., & Kołaczowski, Z. 2016, *MNRAS*, 455, 1728
- Mennickent, R. E., Sterken, C., & Vogt, N. 1997, *A&A*, 326, 1167
- 1400 Merrill, P. W. 1949, *ApJ*, 110, 59
- Mikulášek, Z., Harmanec, P., Grygar, J., & Žďárský, F. 1978, *Bulletin of the Astronomical Institutes of Czechoslovakia*, 29, 44
- Mills, J., J., Snedden, S. A., & Spear, G. G. 1979, *PASP*, 91, 614
- Mourard, D., Brož, M., Nemravová, J. A., et al. 2018, *A&A*, 618, A112
- Mourard, D., Monnier, J. D., Meilland, A., et al. 2015, *A&A*, 577, A51
- Muterspaugh, M. W., Hartkopf, W. I., Lane, B. F., et al. 2010, *AJ*, 140, 1623
- Neiner, C., de Batz, B., Cochard, F., et al. 2011, *AJ*, 142, 149
- Nemravová, J., Harmanec, P., Koubský, P., et al. 2012, *A&A*, 537, A59
- Nemravová, J., Harmanec, P., Kubát, J., et al. 2010, *A&A*, 516, A80
- 1410 Ozaki, A. T. 1991, *PASJ*, 43, 75
- Pavlovski, K. 1979, *A&A*, 76, 362
- Pavlovski, K. & Božić, H. 1982, *Hvar Observatory Bulletin*, 6, 45
- Pavlovski, K., Harmanec, P., Božić, H., et al. 1997, *A&AS*, 125, 75
- Pavlovski, K., Harmanec, P., Horn, J., et al. 1979, *Information Bulletin on Variable Stars*, 1689, 1
- Pearce, J. A. & Hill, G. 1975, *Publications of the Dominion Astrophysical Observatory Victoria*, 14, 319
- Percy, J. R. 1982, in *Bulletin of the American Astronomical Society*, Vol. 14, 879
- 1420 Percy, J. R. 1983, *AJ*, 88, 427
- Percy, J. R., Harlow, J., Hayhoe, K. A. W., et al. 1997, *PASP*, 109, 1215
- Percy, J. R. & Lane, M. C. 1977, *AJ*, 82, 353
- Percy, J. R., Marinova, M. M., Božić, H., & Harmanec, P. 1999, *A&A*, 348, 553

- Perry, C. L., Olsen, E. H., & Crawford, D. L. 1987, *PASP*, 99, 1184
- Perryman, M. A. C. & ESA. 1997, The HIPPARCOS and TYCHO catalogues (Astrometric and photometric star catalogues derived from the ESA Hipparcos Space Astrometry Mission, Publisher: Noordwijk, Netherlands: ESA Publications Division, 1997, Series: ESA SP Series 1200)
- 1430 Peters, G. J. 1971, *ApJ*, 163, L107
- Peters, G. J. 1972, *PASP*, 84, 334
- Peters, G. J. 1983, *PASP*, 95, 311
- Peters, G. J., Pewett, T. D., Gies, D. R., Touhami, Y. N., & Grundstrom, E. D. 2013, *ApJ*, 765, 2
- Peters, G. J., Wang, L., Gies, D. R., & Grundstrom, E. D. 2016, *ApJ*, 828, 47
- Poeckert, R. 1981a, *PASP*, 93, 297
- Poeckert, R. 1981b, *PASP*, 93, 535
- Polidan, R. S. 1976, in *Be and Shell Stars*, ed. A. Slettebak, Vol. 70, 401
- Rast, R. G., Jones, C. E., Carciofi, A. C., et al. 2024, *ApJ*, 968, 30
- Reimann, H. G., Ossenkopf, V., & Beyersdorfer, S. 1992, *A&A*, 265, 360
- 1440 Richards, M. T., Koubský, P., Šimon, V., et al. 2000, *ApJ*, 531, 1003
- Richardson, N. D., Thizy, O., Bjorkman, J. E., et al. 2021, *MNRAS*, 508, 2002
- Roberts, Lewis C., J., Turner, N. H., & ten Brummelaar, T. A. 2007, *AJ*, 133, 545
- Rufener, F. 1986, *A&A*, 165, 275
- Ruždjak, D., Božić, H., Harmanec, P., et al. 2009, *A&A*, 506, 1319
- Saad, S. M., Kubát, J., Hadrava, P., et al. 2005, *Ap&SS*, 296, 173
- Saad, S. M., Kubát, J., Koubský, P., et al. 2004, *A&A*, 419, 607
- Saad, S. M., Nouh, M. I., Shokry, A., & Zead, I. 2021, *Rev. Mexicana Astron. Astrofis.*, 57, 91
- 1450 Saio, H., Cameron, C., Kuschnig, R., et al. 2007, *ApJ*, 654, 544
- Sharov, A. S. & Lyutyj, V. M. 1997, *Astronomy Letters*, 23, 93
- Sharov, A. S. & Lyutyj, V. M. 1972, *Peremennye Zvezdy*, 18, 377
- Shokry, A., Rivinius, T., Mehner, A., et al. 2018, *A&A*, 609, A108
- Sigut, T. A. A. & Ghafourian, N. R. 2023, *ApJ*, 948, 34
- Sigut, T. A. A. & Patel, P. 2013, *ApJ*, 765, 41
- Smith, M. A. & Henry, G. W. 2021, *ApJ*, 915, 13
- Stagg, C. R., Božić, H., Fullerton, A. W., et al. 1988, *MNRAS*, 234, 1021
- Stagg, C. R., Božić, H., Fullerton, A. W., et al. 1985, *JRASC*, 79, 243
- Štefl, S., Harmanec, P., Horn, J., et al. 1990, *BAiCz*, 41, 29
- Sterken, C. 1983, *The Messenger*, 33, 10 1460
- Struve, O. 1944, *ApJ*, 99, 295
- Sudar, D., Harmanec, P., Lehmann, H., et al. 2011, *A&A*, 528, A146
- Tarasov, A. E., Berdyugina, S. V., & Berdyugin, A. V. 1998, *Astronomy Letters*, 24, 316
- Tarasov, A. E. & Tuominen, I. 1987, *Publications of the Astronomical Institute of the Czechoslovak Academy of Sciences*, 5, 127
- Thaller, M. L., Bagnuolo, William G., J., Gies, D. R., & Penny, L. R. 1995, *ApJ*, 448, 878
- Torres, G. 2020, *ApJ*, 901, 91
- Touhami, Y., Gies, D. R., Schaefer, G. H., et al. 2013, *ApJ*, 768, 128 1470
- Tycner, C., Ames, A., Zavala, R. T., et al. 2011, *ApJ*, 729, L5
- Štefl, S., Hummel, W., & Rivinius, T. 2000, *A&A*, 358, 208
- Štefl, S., Rivinius, T., Carciofi, A. C., et al. 2009, *A&A*, 504, 929
- Walker, G. A. H., Kuschnig, R., Matthews, J. M., et al. 2005, *ApJ*, 623, L145
- Walker, G. A. H., Yang, S., & Fahlman, G. G. 1979, *ApJ*, 233, 199
- Wang, L., Gies, D. R., & Peters, G. J. 2018, *ApJ*, 853, 156
- Wang, L., Gies, D. R., Peters, G. J., et al. 2021, *AJ*, 161, 248
- Waters, L. B. F. M., Cote, J., & Pols, O. R. 1991, *A&A*, 250, 437
- Weßmayer, D., Przybilla, N., & Butler, K. 2022, *A&A*, 668, A92
- White, T. R., Pope, B. J. S., Antoci, V., et al. 2017, *MNRAS*, 471, 2882 1480
- Wilson, R. H., J. 1950, *AJ*, 55, 153
- Wolf, M., Harmanec, P., Božić, H., et al. 2021, *A&A*, 647, A97
- Young, A. T., Genet, R. M., Boyd, L. J., et al. 1991, *PASP*, 103, 221

Appendix A: Monitoring the atmospheric extinction at Hvar

Determination of linear extinction coefficients for every night of observations, in which the observed standard stars span a sufficient range of air masses (at least 0.2), was a natural part of the data reduction.

It was deemed useful to present these data, relevant to air pollution at Hvar. Their value exceeds the astronomical interest and can be valuable for broader community, ecology studies in particular.

The seasonal variations, plotted in Fig A.1 and already noted by Koubský & Pavlovski (1982), are clearly seen. During the winter seasons, atmospheric extinction is usually lower as the sea does not evaporate too much. During the summer seasons, the values of the extinction coefficients exhibit a large scatter due to local weather changes but on an average they are much higher than in the winter. Another thing worth noting is that the minimum values of the seasonal changes do not show any annual modulation. Such a behavior has also been found at some other sites, for instance at La Silla (Rufener 1986) or Grossschwabhausen observing station (Reimann et al. 1992)). We note that in Fig A.1 there are some values of extinction coefficients lying below the general slope of the minimum extinction, especially for the U band. These extinction coefficients were usually determined on poor photometric nights or the nights where only few extinction standard stars had been measured. Of course, such nights were never used for all-sky photometry.

In Figure A.2 we present a time plot of all extinction coefficients in U , B and V passbands, recorded at Hvar Observatory over the past fifty years. The scatter of the values is mainly caused by seasonal changes discussed above. Nevertheless, a mild growth of the extinction can be seen in 1975 and 1982. Such increases are often related to large volcanic eruptions anywhere in the world. The first maximum, seen in 1975 may be due to eruption of the El Fuego volcano in Guatemala (October 1974). We warn, however, that during the first observing seasons at Hvar, only a limited number of nights and standard stars suitable for accurate determination of the extinction were available.

The next increase of the extinction appeared at the time of eruption of the El Chiichon volcano (March 1982). The growth of extinction coefficients after this eruption was also recorded at other observatories like La Silla (Rufener 1986; Burki et al. 1995), Jena University Observatory (Reimann et al. 1992), or Lowell Observatory (Lockwood & Thompson 1986). However, the effect of the powerful eruption of the St. Helens volcano (March 1980) was not obviously felt at extinction at Hvar. Needless to say, a very small number of observations was made at that time, however. Also the La Silla observations (Rufener 1986) and observations at the Jena University Observatory (Reimann et al. 1992) did not detected this eruption while at the Lowell Observatory (Lockwood & Thompson 1986), the effect was only marginal. The strong Pinatubo eruption (April 1991) was not covered by Hvar observations because of the intermission due to war in Yugoslavia.

A more detailed analysis of the extinction variations at Hvar and their nature will be published elsewhere. Here we only note that already Pavlovski et al. (1979) pointed out that the influence of the molecular absorption (mainly by O_3) is of little effect in the UBV region. The Rayleigh scattering on air molecules that depends on wavelength and the air pressure shows a small range of variability. The most variable component of the extinction is scattering by aerosols. The Hvar Observatory is a low-altitude station and in the vicinity of the sea and is a subject to signifi-




Fig. A.1: The seasonal variations of the extinction coefficients at Hvar Observatory.

cant weather changes even within one night. Obviously the sea spray plays a major role in the large variations of extinction during the summer seasons. Also occasional forest fires, quite frequent on the Adriatic coast, may have significant influence on the measured extinction.




Fig. A.2: The long-time behaviour of the extinction coefficients at Hvar Observatory over the past fifty years.

Table 4: Hvar standard stars with at least 20 observations secured in 2 or more seasons

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
12 Cas	2011	45307.3–60362.4	117	5.385±0.009	5.397±0.010	5.246±0.012	0.012	-0.151
HR 96	2054	57627.5–58887.3	29	5.746±0.009	5.683±0.009	5.370±0.011	-0.063	-0.313
HR 113	2626	45216.5–60362.3	161	5.940±0.010	5.953±0.011	5.603±0.013	0.013	-0.350
HR 189	4142	45212.6–60583.5	847	5.674±0.008	5.549±0.009	4.978±0.011	-0.125	-0.571
ν 1 Cas	5234	45216.5–59172.4	94	4.842±0.012	6.057±0.010	7.307±0.011	1.215	1.249
HR 285	5848	56639.2–57957.5	62	4.261±0.012	5.481±0.011	6.798±0.012	1.219	1.317
HR 286	5914	54015.4–58385.4	124	6.488±0.010	6.578±0.011	6.652±0.013	0.090	0.074
HR 289	6114	45648.2–60583.5	615	6.472±0.009	6.721±0.009	6.814±0.010	0.248	0.093
HR 333	6798	54015.4–57656.5	35	5.619±0.011	5.641±0.010	5.661±0.013	0.022	0.020
51 And	9927	44956.2–60583.5	925	3.600±0.010	4.881±0.010	6.312±0.011	1.281	1.432
HR 488	10362	54719.5–57779.4	326	6.337±0.009	6.367±0.010	5.942±0.011	0.030	-0.425
2 Per	11291	44956.2–60583.5	754	5.713±0.008	5.646±0.009	5.369±0.011	-0.067	-0.277
HR 561	11875	54719.5–57779.4	257	6.026±0.009	5.996±0.009	5.588±0.010	-0.030	-0.408
47 Cas	12230	55645.4–57615.5	21	5.294±0.014	5.617±0.013	5.658±0.015	0.323	0.041
4 Per	12303	44956.2–60583.5	853	5.009±0.008	4.936±0.009	4.631±0.011	-0.073	-0.305
HR 753	16160	46324.6–59275.3	172	5.815±0.011	6.792±0.010	7.585±0.012	0.977	0.793
HD 16578	16578	52584.3–55068.6	245	9.029±0.022	9.113±0.015	9.186±0.022	0.084	0.073
24 Per	18449	44865.6–59179.4	48	4.959±0.009	6.191±0.011	7.465±0.015	1.232	1.274
HD 18962	18962	42607.5–60323.3	163	7.939±0.011	8.547±0.012	8.644±0.014	0.607	0.098
HD 18992	18992	52584.3–55068.6	190	8.456±0.014	8.545±0.018	8.005±0.018	0.089	-0.540
HD 19193	19193	42607.6–58002.6	109	8.140±0.011	9.396±0.013	10.696±0.022	1.256	1.300
HD 19556	19556	55795.5–60323.3	92	8.441±0.012	9.086±0.011	9.215±0.014	0.645	0.129
ω Per	19656	44861.6–59106.5	169	4.636±0.010	5.748±0.010	6.792±0.014	1.112	1.044
HR 950	19736	44861.6–59097.5	142	6.160±0.011	6.060±0.009	5.486±0.014	-0.099	-0.574
HR 1034	21278	45307.4–59176.4	58	4.999±0.009	4.908±0.012	4.366±0.013	-0.092	-0.541
4 Tau	21686	54116.2–60323.4	390	5.150±0.009	5.105±0.010	5.012±0.012	-0.045	-0.093
HR 1074	21856	44988.3–59179.4	102	5.902±0.010	5.836±0.011	5.005±0.015	-0.066	-0.831
6 Tau	21933	54116.2–60323.4	331	5.774±0.009	5.698±0.010	5.398±0.012	-0.076	-0.300
HR 1133	23193	44988.3–59106.5	49	5.615±0.012	5.675±0.013	5.772±0.016	0.060	0.096
16 Tau	23288	44863.6–60583.5	437	5.476±0.008	5.438±0.009	5.097±0.011	-0.038	-0.341
18 Tau	23324	43140.3–60583.5	621	5.674±0.009	5.601±0.010	5.231±0.011	-0.073	-0.370
43 Per	24546	45307.4–59176.4	30	5.287±0.014	5.704±0.016	5.713±0.018	0.417	0.009
HR 1229	24982	47444.4–59179.4	44	6.506±0.007	6.613±0.009	6.700±0.010	0.107	0.087
HR 1243	25339	56523.6–60583.5	223	5.670±0.011	5.689±0.011	5.267±0.013	0.019	-0.422
ν Tau	25490	41691.2–59248.3	22	3.906±0.014	3.934±0.013	3.999±0.014	0.028	0.064
37 Tau	25604	44863.6–59598.3	240	4.381±0.010	5.440±0.009	6.411±0.010	1.058	0.971
λ Per	25642	48273.3–58888.4	78	4.289±0.006	4.266±0.007	4.258±0.009	-0.023	-0.008
HR 1284	26171	56523.6–60583.5	175	5.954±0.012	5.994±0.011	5.917±0.012	0.040	-0.077
σ 2 Eri	26965	41691.2–59275.3	20	4.426±0.009	5.237±0.011	5.679±0.011	0.812	0.442
HR 1482	29526	48273.3–58888.3	75	5.674±0.005	5.661±0.006	5.630±0.009	-0.013	-0.030
59 Per	29722	44926.5–58889.4	37	5.308±0.009	5.324±0.007	5.351±0.010	0.016	0.027
HD 31894	31894	55574.3–58864.4	49	8.484±0.012	8.529±0.015	7.883±0.016	0.045	-0.645
HR 1644	32655	45307.5–59103.5	205	6.222±0.008	6.667±0.009	7.003±0.012	0.445	0.336
μ Aur	33641	44926.5–58889.4	48	4.826±0.013	5.018±0.022	5.119±0.041	0.192	0.101
HR 1725	34332	55858.6–59179.6	32	6.233±0.010	7.610±0.011	9.247±0.024	1.376	1.638
λ Aur	34411	45307.5–59103.5	260	4.708±0.008	5.317±0.009	5.463±0.011	0.609	0.146
HR 1738	34557	55858.6–59179.6	24	5.478±0.007	5.593±0.007	5.682±0.010	0.115	0.088
ρ Aur	34759	55858.6–59179.6	24	5.223±0.006	5.079±0.007	4.485±0.010	-0.144	-0.594
HD 35619	35619	41691.2–59248.4	37	8.533±0.015	8.802±0.012	8.140±0.014	0.269	-0.662
HD 35653	35653	41691.2–59248.4	36	7.451±0.022	7.586±0.018	6.883±0.013	0.135	-0.703
33 Ori	36351	51512.5–60362.4	249	5.458±0.008	5.274±0.008	4.446±0.009	-0.184	-0.828
HR 1860	36589	43136.4–60583.6	892	6.203±0.009	6.176±0.009	5.779±0.011	-0.028	-0.397
HR 1861	36591	53744.4–60362.4	198	5.341±0.009	5.150±0.009	4.223±0.011	-0.190	-0.927
HR 1871	36741	51512.5–54021.6	65	6.584±0.009	6.398±0.009	5.607±0.010	-0.186	-0.791
121 Tau	36819	44902.6–60583.6	170	5.380±0.010	5.300±0.009	4.668±0.012	-0.080	-0.632
126 Tau	37711	43136.4–60574.6	704	4.847±0.009	4.723±0.010	4.075±0.012	-0.125	-0.648
51 Ori	37984	51512.5–60362.3	126	4.930±0.010	6.078±0.010	7.139±0.012	1.147	1.062
χ 1 Ori	39587	43136.4–60575.6	467	4.406±0.011	4.992±0.011	5.060±0.013	0.586	0.069
HR 2154	41692	44902.6–52561.6	72	5.379±0.012	5.252±0.013	4.707±0.016	-0.127	-0.545

Table 4: continued

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
HR 2205	42690	44902.6–59974.4	369	5.055±0.009	4.860±0.010	4.077±0.012	-0.196	-0.783
HR 2218	43023	44902.7–59974.4	201	5.860±0.009	6.764±0.010	7.401±0.013	0.904	0.636
HR 2248	43526	44993.4–58892.4	67	6.565±0.010	6.450±0.011	5.946±0.015	-0.114	-0.504
HR 2300	44783	44993.4–58892.5	75	6.268±0.011	6.193±0.012	5.904±0.018	-0.076	-0.289
10 Mon	45546	44902.6–59974.4	271	5.055±0.009	4.876±0.009	4.118±0.011	-0.179	-0.758
ψ 4 Aur	47914	45023.4–58925.4	31	5.039±0.010	6.530±0.011	8.372±0.016	1.491	1.842
18 Mon	49293	44993.4–58892.5	54	4.506±0.010	5.605±0.011	6.637±0.016	1.099	1.031
HR 2532	49949	45023.4–58925.4	46	6.263±0.010	6.486±0.009	6.614±0.013	0.223	0.128
16 Lyn	50973	45023.4–58925.4	24	4.919±0.008	4.948±0.009	5.004±0.013	0.030	0.055
HR 2645	52860	45023.4–58925.4	26	6.381±0.009	6.342±0.009	6.091±0.011	-0.038	-0.251
1 CMi	58187	44984.4–59300.4	515	5.383±0.009	5.488±0.009	5.620±0.011	0.104	0.132
ϵ CMi	58367	44984.4–59300.4	273	5.016±0.010	6.013±0.009	6.791±0.012	0.997	0.777
HR 2858	59059	44984.4–59300.4	367	6.247±0.009	6.202±0.009	6.102±0.011	-0.045	-0.100
HD 61341	61341	55586.5–57792.5	31	7.980±0.014	8.175±0.006	8.276±0.025	0.195	0.101
ζ CMi	63975	45056.3–58929.4	47	5.134±0.009	5.016±0.011	4.539±0.011	-0.118	-0.477
HD 65005	65005	52655.4–58946.3	63	8.003±0.009	8.115±0.012	8.247±0.016	0.111	0.133
HD 65199	65199	52655.4–58946.3	81	7.682±0.010	7.600±0.009	7.214±0.012	-0.082	-0.385
HR 3145	66141	53026.5–58929.4	26	4.414±0.008	5.662±0.011	6.949±0.014	1.247	1.287
30 Mon	71155	43143.5–58929.4	45	3.903±0.010	3.887±0.008	3.862±0.010	-0.016	-0.024
39 Cnc	73665	41691.6–59276.4	21	6.427±0.013	7.400±0.010	8.225±0.011	0.973	0.825
42 Cnc	73785	41691.6–59276.4	25	6.840±0.012	7.042±0.013	7.203±0.010	0.202	0.161
HR 3608	77692	55879.6–58946.4	32	6.454±0.006	6.491±0.005	6.573±0.006	0.037	0.081
HD 81772	81772	55879.6–58946.4	43	8.256±0.010	8.570±0.008	8.732±0.012	0.314	0.162
HD 82861	82861	55879.6–58946.4	61	7.074±0.008	7.218±0.008	7.312±0.007	0.144	0.094
σ Leo	83808	46095.5–58946.3	74	3.523±0.012	4.001±0.011	4.211±0.013	0.478	0.210
ν Leo	86360	46095.5–58946.3	61	5.275±0.013	5.222±0.012	5.101±0.014	-0.053	-0.121
λ UMa	89021	43143.6–50902.5	75	3.456±0.012	3.479±0.010	3.558±0.012	0.022	0.079
43 Leo	89962	46095.5–58892.6	39	6.105±0.014	7.204±0.010	8.317±0.021	1.099	1.112
49 UMa	95310	50854.4–50902.5	71	5.082±0.016	5.317±0.010	5.500±0.013	0.236	0.182
51 UMa	95934	50854.4–50902.5	80	6.031±0.016	6.187±0.011	6.288±0.013	0.156	0.101
55 UMa	98353	50854.4–50902.5	63	4.774±0.015	4.881±0.010	4.919±0.015	0.107	0.038
HD 104316	104316	45357.6–58924.5	77	6.775±0.008	6.867±0.008	6.947±0.013	0.092	0.080
HR 4609	104985	44984.7–59733.4	212	5.825±0.013	6.847±0.013	7.693±0.016	1.022	0.846
HR 4687	107193	44984.7–59733.4	359	5.492±0.010	5.534±0.011	5.611±0.013	0.042	0.077
HR 5018	115612	45357.6–59733.4	165	6.220±0.008	6.165±0.008	6.028±0.009	-0.055	-0.137
81 UMa	118214	41828.4–59025.4	30	5.637±0.009	5.608±0.010	5.544±0.009	-0.029	-0.064
HR 5162	119476	52444.4–52445.5	25	5.845±0.009	5.916±0.010	5.983±0.017	0.071	0.067
HD 119581	119581	52444.4–52445.5	23	6.614±0.008	6.738±0.011	6.845±0.016	0.124	0.107
HR 5216	120874	41828.4–59025.4	45	6.490±0.011	6.578±0.012	6.641±0.014	0.089	0.063
50 Boo	136849	45916.3–59032.4	76	5.393±0.009	5.334±0.008	5.143±0.010	-0.059	-0.191
HR 5760	138341	45916.3–59032.4	97	6.469±0.010	6.667±0.010	6.795±0.013	0.199	0.128
BD+64 1078	139549	53491.4–59344.4	178	9.141±0.011	9.546±0.012	9.503±0.017	0.406	-0.044
BD+64 1079	139703	53491.4–59344.4	141	9.312±0.011	10.233±0.011	10.812±0.016	0.921	0.578
HD 141930	141930	45121.4–58564.6	76	7.716±0.008	7.815±0.008	7.916±0.012	0.099	0.102
λ Lib	142096	52068.4–60518.4	100	5.026±0.010	5.035±0.011	4.430±0.014	0.009	-0.605
χ Her	142373	42994.3–60467.4	472	4.606±0.008	5.176±0.009	5.185±0.011	0.569	0.009
λ CrB	142908	44432.3–59032.4	129	5.436±0.010	5.781±0.011	5.816±0.011	0.345	0.036
ν Her	144206	41528.5–60467.4	720	4.736±0.008	4.642±0.009	4.315±0.011	-0.094	-0.327
ω 1 Sco	144470	42241.4–60518.4	121	3.935±0.010	3.932±0.012	3.092±0.013	-0.003	-0.841
φ Her	145389	41536.4–60467.4	544	4.249±0.007	4.189±0.009	3.937±0.010	-0.061	-0.251
HR 6096	147550	52494.3–54296.4	80	6.243±0.013	6.301±0.010	6.293±0.013	0.058	-0.009
ρ Oph	147933	52494.3–56094.4	21	4.567±0.031	4.802±0.030	4.227±0.031	0.235	-0.575
ν Oph	148357	52494.3–56094.4	135	4.636±0.010	4.814±0.013	4.888±0.013	0.178	0.074
16 Oph	151133	52494.3–54280.5	50	6.037±0.007	6.017±0.009	5.878±0.011	-0.020	-0.139
HR 6278	152585	52494.3–53192.4	32	6.584±0.012	6.715±0.015	6.829±0.019	0.130	0.114
ϵ Her	153808	52068.3–55120.3	95	3.918±0.010	3.895±0.010	3.804±0.011	-0.023	-0.091
HR 6353	154445	54275.4–58002.3	90	5.623±0.010	5.795±0.012	5.171±0.016	0.172	-0.623
HR 6361	154660	54275.4–58006.3	363	6.360±0.010	6.572±0.011	6.667±0.014	0.212	0.095
HR 6367	154895	54275.4–58002.3	275	6.061±0.009	6.140±0.010	6.159±0.011	0.078	0.019
72 Her	157214	52068.3–55120.3	95	5.385±0.011	6.005±0.010	6.090±0.011	0.620	0.085

Table 4: continued

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
77 Her	158414	43313.4–60574.3	814	5.851±0.008	5.979±0.009	6.118±0.010	0.128	0.139
30 Dra	162579	41536.4–60574.3	710	5.040±0.008	5.073±0.009	5.080±0.010	0.033	0.007
ξ Dra	163588	42241.4–60574.3	626	3.745±0.010	4.925±0.010	6.150±0.011	1.180	1.225
HR 6689	163624	43712.4–59043.5	77	5.953±0.012	6.078±0.012	6.167±0.017	0.125	0.089
HR 6690	163641	45105.5–60535.3	146	6.290±0.009	6.296±0.010	6.062±0.011	0.006	-0.234
93 Her	164349	44799.3–60535.4	356	4.680±0.009	5.923±0.009	7.141±0.010	1.243	1.219
HR 6719	164432	45105.5–60535.4	176	6.352±0.009	6.277±0.011	5.520±0.012	-0.075	-0.756
71 Oph	165760	45105.5–59793.4	106	4.663±0.008	5.615±0.009	6.356±0.011	0.952	0.741
102 Her	166182	44799.4–60535.4	491	4.359±0.008	4.189±0.010	3.392±0.011	-0.170	-0.797
101 Her	166230	46269.4–60531.4	343	5.115±0.008	5.278±0.009	5.458±0.010	0.162	0.180
15 Sgr	167264	41824.6–59027.5	32	5.315±0.018	5.388±0.018	4.524±0.020	0.073	-0.864
21 Sgr	169420	41824.6–59027.5	41	4.801±0.016	6.122±0.020	7.106±0.025	1.321	0.984
HR 6900	169578	44433.4–59044.5	271	6.732±0.010	6.758±0.011	5.561±0.013	0.026	-0.197
HR 6925	170137	44433.4–59044.5	145	6.003±0.012	7.659±0.012	9.520±0.019	1.656	1.860
HR 6928	170200	44433.4–59044.5	313	5.711±0.009	5.687±0.010	5.329±0.014	-0.024	-0.358
HR 6997	172044	49538.5–56899.4	58	5.413±0.006	5.319±0.009	4.805±0.014	-0.094	-0.514
HR 7028	172883	42993.5–59043.5	726	6.015±0.009	5.942±0.009	5.731±0.012	-0.072	-0.211
HR 7060	173664	42993.4–59051.5	891	6.207±0.009	6.282±0.009	6.434±0.013	0.074	0.153
ν2 Lyr	174602	49538.5–57973.4	333	5.240±0.008	5.338±0.011	5.438±0.011	0.098	0.100
HR 7123	175225	42998.4–59051.5	492	5.523±0.009	6.357±0.010	6.878±0.014	0.835	0.520
HD 175406	175406	55357.5–56541.4	292	8.133±0.008	8.395±0.008	8.480±0.011	0.261	0.086
HD 175469	175469	55357.5–56541.4	268	8.995±0.010	9.216±0.009	9.306±0.011	0.221	0.090
γ Lyr	176437	46969.4–59180.2	431	3.252±0.009	3.184±0.013	3.148±0.013	-0.068	-0.036
HR 7239	177817	53584.4–59098.3	75	6.012±0.010	5.993±0.010	5.648±0.010	-0.019	-0.345
51 Dra	178201	42994.4–56094.4	65	5.419±0.018	5.395±0.023	5.328±0.030	-0.024	-0.067
HD 180889	180889	46238.5–47374.4	157	6.907±0.008	7.078±0.010	7.239±0.015	0.172	0.161
HD 181751	181751	46238.5–47374.4	108	6.571±0.008	6.500±0.011	6.150±0.015	-0.070	-0.351
HR 7378	182645	53936.5–59098.3	46	5.716±0.010	5.753±0.012	5.424±0.011	0.037	-0.329
HR 7379	182678	54280.4–59098.3	45	6.719±0.009	6.766±0.008	6.695±0.011	0.047	-0.072
HR 7397	183227	44079.5–60575.4	1566	5.844±0.009	5.861±0.010	5.482±0.013	0.017	-0.379
HR 7404	183387	44079.5–60575.4	964	6.251±0.009	7.570±0.011	8.931±0.015	1.319	1.361
9 Vul	184606	46679.4–59025.5	103	5.008±0.009	4.909±0.010	4.499±0.012	-0.099	-0.410
HR 7438	184663	44079.5–60575.4	1137	6.373±0.009	6.782±0.010	6.748±0.012	0.409	-0.034
β Sge	185958	46238.5–58690.6	111	4.392±0.008	5.426±0.010	6.328±0.016	1.034	0.902
HD 185978	185978	54751.2–58402.4	102	7.892±0.011	8.398±0.014	8.493±0.016	0.506	0.095
HD 186239	186239	58324.5–58402.4	75	7.362±0.011	7.583±0.011	7.675±0.014	0.221	0.092
HR 7512	186568	45900.5–58696.5	597	6.077±0.009	6.003±0.009	5.713±0.011	-0.074	0.290
15 Cyg	186675	46679.4–60575.4	88	4.900±0.008	5.837±0.012	6.540±0.013	0.936	0.704
γ Aql	186791	46969.4–47061.3	20	2.717±0.010	4.231±0.013	5.945±0.020	1.515	1.713
17 Cyg	187013	57210.5–58696.5	81	5.007±0.010	5.469±0.012	5.471±0.013	0.462	0.001
HR 7550	187458	45900.5–58696.5	722	6.659±0.009	7.083±0.010	7.031±0.012	0.424	-0.052
HD 187664	187664	54280.5–59082.3	39	6.890±0.011	7.204±0.012	7.435±0.012	0.314	0.232
HD 188170	188170	43027.4–58682.6	104	7.352±0.013	7.275±0.015	6.900±0.017	-0.077	-0.375
13 Vul	188260	46665.5–59025.5	160	4.585±0.008	4.536±0.009	4.403±0.012	-0.049	-0.133
β Aql	188512	42654.3–57970.5	21	3.722±0.015	4.576±0.015	5.064±0.012	0.854	0.488
22 Cyg	188892	46238.5–60583.3	362	4.953±0.008	4.862±0.010	4.344±0.013	-0.090	-0.518
17 Vul	190993	44817.4–60583.3	225	5.082±0.010	4.908±0.011	4.228±0.013	-0.174	-0.680
18 Vul	191747	44432.5–60583.3	178	5.529±0.009	5.603±0.009	5.722±0.012	0.074	0.119
19 Vul	192004	44432.5–60583.3	118	5.492±0.012	6.907±0.012	8.406±0.016	1.415	1.499
33 Cyg	192696	46323.3–59101.4	59	4.289±0.010	4.398±0.012	4.501±0.014	0.109	0.103
36 Cyg	193369	46238.5–60583.3	273	5.593±0.009	5.633±0.009	5.663±0.013	0.040	0.030
HR 7815	194668	44433.5–59087.6	355	6.511±0.009	6.496±0.011	6.447±0.014	-0.015	-0.049
42 Cyg	195324	46680.4–59077.5	40	5.874±0.010	6.321±0.013	6.531±0.027	0.447	0.210
HD 197618	197618	53578.3–57964.5	127	6.995±0.009	7.154±0.009	7.241±0.011	0.159	0.087
η Cep	198149	41528.4–59082.5	296	3.426±0.010	4.337±0.011	4.953±0.015	0.911	0.616
HD 198679	198679	53686.2–57966.4	23	6.905±0.011	6.852±0.012	6.694±0.015	-0.053	-0.157
HD 198692	198692	51426.4–59101.4	65	6.660±0.011	7.672±0.010	8.483±0.017	1.012	0.811
HR 7996	198820	52165.4–59101.4	35	6.436±0.010	6.299±0.007	5.669±0.009	-0.138	-0.630
HD 199007	199007	51402.4–59101.4	154	7.958±0.012	7.907±0.009	7.587±0.011	-0.051	-0.320
HD 199311	199311	46269.5–60583.4	841	6.687±0.008	6.765±0.009	6.857±0.011	0.078	0.092

Table 4: continued

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
HD 199479	199479	46269.5–52094.5	135	6.845±0.010	6.804±0.009	6.587±0.013	-0.041	-0.217
HIP 103770	200270	52500.4–52549.4	163	7.768±0.010	8.227±0.012	8.171±0.013	0.459	-0.057
HIP 103874	200468	52500.4–52549.4	142	7.862±0.009	8.039±0.012	8.157±0.013	0.177	0.119
HR 8119	202214	45538.5–59101.4	130	5.617±0.007	5.737±0.008	4.982±0.010	0.120	-0.755
HD 202349	202349	44817.4–59082.5	217	7.362±0.010	7.195±0.009	6.222±0.012	-0.167	-0.974
HR 8161	203245	51067.3–60583.4	937	5.762±0.008	5.633±0.009	5.115±0.010	-0.128	-0.519
α Cep	203280	46988.4–52494.5	24	2.457±0.010	2.676±0.013	2.786±0.015	0.219	0.110
70 Cyg	204403	44817.4–59082.5	163	5.315±0.007	5.166±0.009	4.505±0.012	-0.149	-0.661
72 Cyg	205512	44817.4–60531.5	652	4.891±0.009	5.981±0.010	6.991±0.011	1.090	1.011
HD 206259	206259	45155.5–59097.4	32	7.534±0.010	7.596±0.011	7.032±0.014	0.062	-0.563
HD 206694	206694	57581.5–57996.5	157	8.804±0.012	9.319±0.011	9.393±0.014	0.515	0.074
π 2 Cyg	207330	42244.5–59097.4	83	4.244±0.008	4.116±0.011	3.404±0.015	-0.128	-0.713
12 Cep	207528	58740.4–59043.6	24	5.517±0.012	7.086±0.011	9.021±0.024	1.569	1.935
HR 8342	207636	57581.5–58006.5	168	6.475±0.009	6.466±0.011	6.437±0.012	-0.009	-0.029
HD 207793	207793	45155.5–59087.5	21	6.560±0.010	6.940±0.011	6.424±0.012	0.380	-0.516
HR 8473	210873	57581.5–58006.5	193	6.389±0.011	6.327±0.011	6.161±0.012	-0.063	-0.165
2 Lac	212120	41536.5–59180.2	88	4.550±0.011	4.445±0.013	3.916±0.014	-0.105	-0.529
β Lac	212496	45155.5–59097.4	40	4.448±0.010	5.458±0.012	6.249±0.016	1.009	0.791
4 Lac	212593	41536.5–58740.5	169	4.578±0.011	4.666±0.012	4.314±0.014	0.089	-0.352
6 Lac	213420	41929.5–59180.3	31	4.509±0.011	4.421±0.011	3.691±0.012	-0.088	-0.730
α Lac	213558	46280.5–50341.5	57	3.775±0.010	3.784±0.011	3.802±0.011	0.009	0.018
9 Lac	214454	42996.5–50341.5	44	4.641±0.011	4.887±0.010	5.005±0.012	0.246	0.118
15 Lac	216397	42993.5–46696.5	54	4.921±0.023	6.536±0.038	8.488±0.034	1.615	1.952
HR 8708	216608	46280.5–46696.5	25	5.771±0.009	6.060±0.013	6.170±0.016	0.290	0.110
HR 8733	217101	44432.5–60583.4	1455	6.175±0.009	6.028±0.009	5.220±0.011	-0.147	-0.808
2 And	217782	42993.5–60583.4	949	5.108±0.009	5.194±0.010	5.304±0.011	0.086	0.110
3 And	218031	46664.6–60189.5	263	4.662±0.008	5.718±0.010	6.618±0.012	1.056	0.899
1 Cas	218376	50854.2–51080.5	40	4.828±0.018	4.792±0.013	3.947±0.027	-0.035	-0.845
4 And	218452	43140.2–60583.4	892	5.301±0.011	6.724±0.011	8.443±0.015	1.423	1.720
5 And	218470	42266.5–60197.6	1203	5.686±0.009	6.116±0.010	6.100±0.012	0.430	-0.016
HR 8806	218525	43712.5–59077.6	25	6.524±0.011	6.692±0.010	6.892±0.010	0.168	0.200
7 And	219080	45561.5–60197.6	564	4.535±0.008	4.828±0.010	4.861±0.011	0.293	0.033
HD 219326	219326	53241.5–58006.5	318	8.361±0.010	8.462±0.011	8.575±0.012	0.101	0.114
HR 8870	219891	43712.5–59077.6	54	6.528±0.008	6.689±0.009	6.827±0.011	0.161	0.138
10 And	219981	46664.5–51066.6	25	5.788±0.012	7.321±0.015	9.185±0.019	1.533	1.865
HD 220073	220073	52136.6–58006.5	454	7.699±0.009	8.119±0.010	8.122±0.012	0.420	0.003
ι And	222173	42266.5–51372.4	25	4.304±0.016	4.205±0.013	3.886±0.021	-0.099	-0.319
κ And	222439	42266.5–60583.4	406	4.155±0.009	4.068±0.010	3.849±0.011	-0.086	-0.220
HR 9011	223229	45297.2–60583.4	395	6.085±0.009	5.957±0.008	5.289±0.010	-0.127	-0.668
HR 9057	224342	46287.6–58716.4	56	6.019±0.010	6.739±0.015	7.067±0.021	0.720	0.328
σ Cas	224572	50854.2–51080.5	38	4.880±0.013	4.805±0.010	4.006±0.016	-0.075	-0.799
HD 224890	224890	54754.3–55120.6	34	6.510±0.015	6.694±0.017	6.791±0.019	0.184	0.098
BD+52 2289	–	42994.4–54294.4	32	9.761±0.017	10.201±0.051	10.199±0.059	0.440	-0.002
BD+52 2288	–	42994.4–54294.4	45	9.029±0.018	9.490 ±.021	9.441±0.033	0.461	-0.048

Table 5: Less frequently observed standard stars

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
HR 1	3	46287.6–46290.6	9	6.718±0.008	6.783±0.011	6.858±0.014	0.065	0.075
HD 567	567	57634.5–58887.3	17	7.221±0.017	7.191±0.024	6.801±0.023	-0.029	-0.390
χ Peg	1013	42266.5–42278.5	3	4.866±0.011	6.504±0.019	8.350±0.017	1.638	1.846
HD 1687	1687	56092.5–56092.5	2	8.138±0.003	9.052±0.007	9.648±0.012	0.914	0.596
BD+81 13/A	3440	46689.6–46696.6	2	6.367±0.009	6.921±0.015	6.905±0.022	0.553	-0.016
ν And	4727	43136.2–59179.4	18	4.541±0.011	4.399±0.009	3.812±0.010	-0.142	-0.587
20 Cet	5112	43140.2–43152.3	2	4.738±0.059	6.351±0.023	8.287±0.030	1.613	1.936
ν 2 Cas	5395	52488.5–54021.5	16	4.639±0.007	5.597±0.008	6.266±0.012	0.958	0.669
BD+81 27/G	5817	46689.6–46689.6	1	8.429±0.000	9.019±0.000	9.015±0.000	0.590	-0.004
BD+81 29/F	5905	46689.6–46689.6	1	8.519±0.000	9.163±0.000	9.336±0.000	0.644	0.173

Table 5: continued

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
BD+81 30/E	6006	46689.6–46690.6	2	7.968±0.026	7.979±0.028	7.904±0.035	0.012	-0.075
80 Psc	6763	43140.2–43152.3	2	5.514±0.027	5.852±0.001	5.860±0.024	0.338	0.008
BD+80 35/C	7471	46689.6–46689.6	1	7.209±0.000	7.568±0.000	7.645±0.000	0.359	0.077
BD+80 36/B	7505	46689.6–46689.6	1	6.688±0.000	6.715±0.000	6.729±0.000	0.027	0.014
ξ And	8207	43136.3–43152.3	4	4.880±0.013	5.985±0.019	7.004±0.019	1.104	1.019
ψ Cas	8491	45216.5–45216.5	1	4.737±0.000	5.793±0.000	6.734±0.000	1.056	0.941
ν And	9826	46324.6–46324.6	3	4.094±0.002	4.623±0.002	4.691±0.003	0.529	0.068
τ And	10205	43136.3–43152.3	4	4.970±0.028	4.878±0.013	4.480±0.019	-0.091	-0.398
107 Psc	10476	46324.6–46324.6	3	5.267±0.007	6.097±0.009	6.604±0.019	0.830	0.507
HR 502	10857	51447.5–51928.3	9	6.256±0.008	6.323±0.006	6.406±0.008	0.067	0.083
HR 540	11408	51435.6–51928.3	12	6.449±0.005	6.638±0.006	6.751±0.012	0.188	0.113
HR 572	12005	56898.4–56898.4	2	6.056±0.004	7.195±0.003	8.033±0.006	1.140	0.837
50 Cas	12216	47371.6–47373.6	2	3.991±0.021	4.010±0.028	4.053±0.038	0.019	0.042
HD 13841	13841	46679.6–46680.6	7	7.365±0.014	7.606±0.009	6.977±0.009	0.242	-0.629
HD 14369	14369	54018.5–56095.4	19	8.097±0.012	8.499±0.015	8.440±0.016	0.402	-0.059
π Cet	17081	43136.4–43152.3	4	4.232±0.023	4.149±0.023	3.724±0.025	-0.083	-0.425
HR 975	20193	54026.5–54026.6	4	6.311±0.005	6.668±0.008	6.642±0.007	0.357	-0.025
α Per	20902	47049.5–52494.5	10	1.798±0.011	2.271±0.010	2.655±0.025	0.473	0.383
HR 1019	20995	54026.5–54026.6	4	5.795±0.009	5.780±0.006	5.623±0.004	-0.015	-0.157
HR 1041	21402	54026.5–54026.6	4	5.733±0.002	5.769±0.003	5.836±0.007	0.036	0.067
27 Tau	23850	52946.5–56508.6	11	3.642±0.008	3.548±0.014	3.196±0.011	-0.093	-0.352
HR 1188	23985	43140.3–43152.3	13	5.263±0.031	5.480±0.033	5.562±0.032	0.217	0.083
HR 1226	24843	52488.5–52488.5	2	6.302±0.005	7.368±0.001	8.273±0.008	1.066	0.905
HR 1234	25152	52502.5–55454.5	2	6.421±0.003	6.411±0.036	6.328±0.023	-0.010	-0.084
HD 25539	25539	55454.5–55454.5	8	6.872±0.017	6.916±0.030	6.323±0.034	0.044	-0.593
π3 Ori	30652	56540.6–57996.6	15	3.186±0.009	3.631±0.014	3.625±0.014	0.446	-0.007
π4 Ori	30836	56540.6–57634.6	10	3.682±0.011	3.513±0.018	2.704±0.010	-0.169	-0.809
HD 31617	31617	56002.3–56002.3	3	7.423±0.011	7.432±0.008	6.692±0.007	0.008	-0.740
HR 1617	32249	51445.6–51520.5	46	4.795±0.008	4.617±0.014	3.876±0.016	-0.178	-0.741
HR 1621	32309	43136.4–43152.3	4	4.902±0.018	4.858±0.032	4.726±0.026	-0.045	-0.131
HD 32328	32328	56002.3–56002.3	3	7.664±0.008	7.616±0.006	7.278±0.007	-0.049	-0.338
HR 1671	33224	51445.6–51520.5	25	5.820±0.015	5.735±0.015	5.344±0.017	-0.086	-0.391
HR 1697	33833	51445.6–51520.5	12	5.928±0.009	6.869±0.011	7.603±0.012	0.941	0.734
α Aur	34029	47049.5–47058.5	4	0.046±0.017	0.864±0.036	1.312±0.039	0.818	0.448
16 Cam	34787	45309.6–45697.4	7	5.260±0.007	5.239±0.015	5.195±0.013	-0.021	-0.044
β Tau	35497	47052.6–47058.5	2	1.660±0.015	1.516±0.003	1.027±0.004	-0.144	-0.489
HR 1833	36166	51943.3–53761.3	19	5.770±0.008	5.575±0.011	4.729±0.009	-0.195	-0.846
38 Ori	36777	51943.3–51943.4	3	5.340±0.011	5.397±0.010	5.448±0.013	0.057	0.051
φ1 Ori	36822	53388.4–53388.4	3	4.388±0.007	4.246±0.002	3.301±0.007	-0.142	-0.945
σ Ori	37468	58499.4–58499.4	1	3.821±0.000	3.582±0.000	2.566±0.000	-0.239	-1.016
ξ Aur	39283	45308.5–55124.5	18	4.983±0.009	5.026±0.014	5.119±0.016	0.043	0.093
δ Aur	40035	45308.5–55124.5	12	3.758±0.009	4.755±0.016	5.587±0.014	0.997	0.832
HR 2224	43157	45359.3–45360.3	5	5.834±0.006	5.668±0.009	4.991±0.007	-0.166	-0.677
HR 2234	43319	45359.3–45360.3	5	6.009±0.013	6.092±0.009	6.172±0.008	0.084	0.080
HD 46149	46149	53388.5–53388.5	3	7.582±0.001	7.753±0.002	6.995±0.007	0.171	-0.758
ψ2 Aur	47174	43140.5–43152.5	3	4.802±0.035	6.058±0.012	7.338±0.039	1.256	1.280
ν3 CMa	47442	43140.5–43152.5	3	4.422±0.030	5.547±0.023	6.553±0.044	1.125	1.006
HD 50109	50109	54862.4–54862.4	4	9.085±0.019	9.079±0.011	8.983±0.017	-0.006	-0.096
HD 50169	50169	54862.4–54862.4	3	9.021±0.008	9.023±0.007	9.018±0.007	0.002	-0.005
HD 50346	50346	54862.4–54862.4	3	8.756±0.009	8.680±0.007	8.107±0.007	-0.076	-0.573
HD 50669	50669	54862.4–54862.4	3	8.929±0.011	9.625±0.013	9.915±0.008	0.696	0.290
HR 2572	50747	54862.4–54862.4	3	5.463±0.004	5.616±0.011	5.788±0.006	0.153	0.172
HD 53035	53035	52940.6–52940.6	13	7.897±0.016	7.827±0.005	7.311±0.037	-0.070	-0.515
λ Gem	56537	45065.3–50865.4	9	3.583±0.008	3.689±0.006	3.818±0.015	0.106	0.129
16 Pup	67797	43140.6–43152.5	3	4.395±0.005	4.249±0.007	3.653±0.017	-0.146	-0.596
9 Hya	74137	43140.6–43152.5	4	4.882±0.011	5.929±0.018	6.845±0.028	1.047	0.916
ι UMa	76644	43143.5–43143.5	1	3.141±0.000	3.351±0.000	3.411±0.000	0.210	0.060
κ UMa	77327	43140.6–43152.5	3	3.580±0.006	3.562±0.008	3.596±0.006	-0.017	0.034
θ Hya	79469	43143.5–43143.5	1	3.928±0.000	3.851±0.000	3.728±0.000	-0.077	-0.123
HR 3751	81817	56747.3–56747.4	4	4.278±0.008	5.756±0.007	7.470±0.005	1.479	1.714

Table 5: continued

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
21 LMi	87696	43143.6–43143.6	1	4.481±0.000	4.687±0.000	4.738±0.000	0.206	0.051
α Leo	87901	46095.5–58941.5	15	1.341±0.010	1.180±0.022	0.851±0.012	-0.161	-0.329
56 UMa	98839	41824.4–41828.3	2	4.997±0.018	5.970±0.033	6.746±0.014	0.972	0.776
γ Crt	99211	41824.4–41828.3	2	3.980±0.038	4.190±0.038	4.197±0.042	0.210	0.007
HD 100972	100972	41824.4–41828.4	2	6.842±0.002	6.838±0.015	6.843±0.002	-0.004	0.004
ζ Crt	102070	41824.4–41828.3	2	4.659±0.018	5.625±0.004	6.311±0.030	0.967	0.685
γ UMa	103287	43143.6–43143.6	1	2.452±0.000	2.465±0.000	2.475±0.000	0.013	0.010
HR 4561	103498	55357.4–55364.4	8	7.031±0.009	7.053±0.010	7.062±0.009	0.021	0.009
HR 4572	103799	55357.4–55364.4	6	6.613±0.009	7.096±0.011	7.082±0.011	0.483	-0.014
HD 103984	103984	55357.4–55364.4	6	6.931±0.009	7.334±0.008	7.319±0.009	0.403	-0.015
δ UMa	166205	46969.4–47006.3	6	3.312±0.006	3.388±0.013	3.445±0.017	0.076	0.057
η Vir	107259	41691.6–41696.6	4	3.861±0.019	3.867±0.027	3.910±0.040	0.006	0.043
6 Dra	109551	51376.4–51379.4	5	4.945±0.010	6.278±0.008	7.488±0.014	1.333	1.209
χ Vir	110014	41691.6–41696.6	4	4.607±0.024	5.794±0.023	7.166±0.042	1.188	1.371
β Com	114710	42646.3–42654.3	4	4.261±0.022	4.843±0.033	4.899±0.024	0.582	0.056
η UMa	120315	46969.4–47006.3	9	1.856±0.010	1.668±0.008	0.995±0.013	-0.189	-0.672
86 UMa	121409	41828.4–59025.4	18	5.740±0.009	5.701±0.012	5.629±0.011	-0.039	-0.072
ϵ Boo	129989	52494.4–52494.4	1	2.362±0.000	3.333±0.000	4.064±0.000	0.971	0.731
χ Boo	135502	52412.4–52500.4	12	5.294±0.007	5.347±0.007	5.418±0.006	0.053	0.071
γ CrB	140436	52412.4–52500.4	12	3.826±0.006	3.827±0.007	3.807±0.005	0.001	-0.020
ϑ Lib	142198	41824.4–58946.6	12	4.165±0.011	5.180±0.013	5.978±0.019	1.014	0.799
ν Sco AB	145502	41824.4–42929.4	7	3.998±0.008	4.047±0.009	3.389±0.012	0.049	-0.658
ψ Oph	147700	42241.4–42244.4	2	4.439±0.016	5.450±0.001	6.283±0.008	1.010	0.834
59 Her	154029	52068.3–55120.3	9	5.278±0.010	5.298±0.007	5.342±0.016	0.020	0.044
α Oph	159561	46969.4–47006.4	11	2.079±0.008	2.227±0.017	2.349±0.017	0.149	0.122
γ Dra	164058	46969.4–47061.3	22	2.227±0.008	3.762±0.010	5.620±0.016	1.535	1.858
HR 6757	165462	43712.4–43722.4	8	6.394±0.012	7.435±0.018	8.255±0.017	1.042	0.819
HD 166101	166101	57652.3–57652.4	4	8.793±0.027	9.574±0.016	9.798±0.042	0.780	0.224
δ UMi	166205	56858.5–56858.5	3	4.363±0.004	4.377±0.003	4.417±0.004	0.014	0.039
HD 166361	166361	57652.3–57652.4	3	7.454±0.018	9.292±0.015	11.089±0.153	1.839	1.796
HD 168443	168443	57652.3–57652.4	4	6.893±0.005	7.634±0.010	7.924±0.016	0.741	0.289
HR 6943	170650	55121.2–55121.3	13	5.877±0.009	5.788±0.015	5.270±0.016	-0.089	-0.518
HR 6966	171245	55121.2–55121.2	2	5.884±0.004	7.365±0.011	9.118±0.022	1.481	1.753
BD+24 3545	174369	55124.3–55124.3	2	6.750±0.014	6.833±0.001	6.898±0.005	0.083	0.066
ξ 2 Sgr	175775	42241.4–53937.5	7	3.537±0.017	4.698±0.024	5.819±0.018	1.162	1.121
λ Lyr	176679	56491.4–56639.2	2	4.961±0.001	6.414±0.018	8.082±0.012	1.454	1.668
ζ Aql	177724	46969.4–47061.3	18	2.990±0.009	2.996±0.011	3.002±0.014	0.006	0.006
β 1 Cyg	183912	46689.3–46689.4	2	3.067±0.003	4.189±0.001	4.864±0.007	1.123	0.674
κ Aql	184915	42647.3–42654.4	3	4.938±0.017	4.925±0.015	4.069±0.016	-0.013	-0.855
HD 187411	187411	57615.4–59082.4	13	8.105±0.009	8.471±0.010	8.601±0.013	0.366	0.130
23 Vul	192806	43027.4–50341.5	13	4.511±0.009	5.780±0.007	6.920±0.015	1.268	1.140
39 Cyg	194317	46688.4–46695.4	4	4.431±0.024	5.781±0.017	7.297±0.019	1.351	1.516
13 Cap	196348	43712.4–43722.5	2	6.791±0.011	8.017±0.008	9.292±0.012	1.227	1.274
71 Aql	196574	41528.4–59179.3	13	4.337±0.009	5.270±0.008	5.954±0.012	0.933	0.684
HR 7922	197226	47017.4–47022.4	7	6.521±0.006	6.396±0.008	5.907±0.017	-0.125	-0.489
ϵ Aqr	198001	41528.4–41528.4	2	3.775±0.022	3.779±0.011	3.790±0.037	0.003	0.012
HD 198424	198424	47017.4–47022.4	12	7.510±0.008	7.410±0.012	6.980±0.020	-0.100	-0.431
56 Cyg	198639	51715.4–51715.4	3	5.068±0.005	5.266±0.010	5.370±0.012	0.198	0.104
HD 198793	198793	53686.2–53686.2	3	7.141±0.003	7.058±0.005	6.592±0.007	-0.082	-0.466
HR 8029	199661	53686.2–53686.2	3	6.237±0.004	6.075±0.006	5.350±0.003	-0.162	-0.725
HD 199781	199781	51376.3–51380.5	5	8.335±0.026	8.789±0.041	8.778±0.075	0.453	-0.010
HD 199890	199890	51715.4–51715.4	3	7.518±0.001	7.422±0.007	7.044±0.012	-0.097	-0.378
HD 199986	199986	51377.4–58351.5	8	7.052±0.038	7.488±0.410	7.755±0.721	0.436	0.267
HR 8049	200205	46691.4–46691.4	2	5.537±0.004	6.964±0.004	8.562±0.007	1.427	1.599
ϑ Cap	200761	42607.5–43722.5	2	4.062±0.005	4.057±0.010	4.058±0.016	-0.005	0.001
HR 8107	201836	43027.4–43027.4	6	6.452±0.007	6.445±0.014	6.064±0.015	-0.007	-0.381
HR 8136	202654	43027.4–43027.4	5	6.445±0.014	6.308±0.009	5.626±0.011	-0.137	-0.682
1 Peg	203504	42237.4–42237.4	1	4.104±0.000	5.195±0.000	6.250±0.000	1.091	1.055
HR 8185	203644	51371.4–51371.5	4	5.688±0.005	6.787±0.015	7.825±0.019	1.099	1.038
κ Cap	206453	42607.6–43713.6	3	4.716±0.024	5.581±0.022	6.113±0.021	0.864	0.532

Table 5: continued

Name of Star	HD or BD number	JDmin - JDmax -2400000	Number of observations	<i>V</i> (mag.)	<i>B</i> (mag.)	<i>U</i> (mag.)	<i>B</i> - <i>V</i> (mag.)	<i>U</i> - <i>B</i> (mag.)
HD 209932	209932	55452.3–55460.5	52	6.522±0.012	6.490±0.015	6.350±0.016	-0.032	-0.140
BD+48 3613	210119	52940.2–52940.4	14	8.328±0.009	8.426±0.009	8.505±0.016	0.098	0.078
HD 210387	210387	55452.3–55460.5	33	6.799±0.013	6.761±0.014	6.486±0.016	-0.038	-0.275
35 Peg	212943	42266.4–42278.4	3	4.765±0.010	5.811±0.022	6.682±0.023	1.046	0.871
HR 8606	214240	43143.2–43326.6	7	6.276±0.016	6.227±0.012	5.664±0.010	-0.049	-0.563
11 Lac	214868	42607.6–42609.6	2	4.492±0.006	5.828±0.001	7.287±0.003	1.335	1.459
HR 8654	215359	58700.6–58700.6	3	5.921±0.008	7.442±0.004	9.139±0.015	1.521	1.698
α Peg	218045	46988.4–47061.3	23	2.482±0.009	2.441±0.011	2.389±0.014	-0.041	-0.052
HD 219430	219430	42278.5–42278.5	1	4.165±0.000	5.262±0.000	6.287±0.000	1.097	1.025
ψ 1 Aqr	219449	42237.6–42237.6	1	4.268±0.000	5.403±0.000	6.415±0.000	1.135	1.012
HR 8857	219668	43712.5–43722.5	5	6.505±0.006	7.547±0.016	8.539±0.014	1.042	0.991
72 Peg	221673	42237.6–42237.6	1	5.010±0.000	6.390±0.000	8.007±0.000	1.380	1.617
ω 2 Aqr	222661	42607.6–43713.6	3	4.454±0.022	4.385±0.024	4.248±0.032	-0.069	-0.137
BD+37 3873	228897	55121.3–55124.4	13	8.923±0.008	9.060±0.009	8.590±0.013	0.137	-0.470
HD 277197	277197	55851.5–55858.7	11	9.490±0.013	9.557±0.015	9.018±0.027	0.067	-0.539
HD 339764	339764	55454.4–55454.4	1	9.599±0.000	9.688±0.000	9.779±0.000	0.089	0.091
ADS 10347B	–	54275.4–54294.4	15	9.630±0.062	10.267±0.097	10.407±0.113	0.637	0.140
ADS 10347AB	–	54281.5–57948.4	15	6.334±0.033	6.581±0.068	6.702±0.131	0.247	0.121
BD+43 1166C	–	55596.4–56002.3	5	10.720±0.338	11.402±0.423	11.712±0.393	0.682	0.310
BD+67 248	–	55851.4–55858.6	11	9.024±0.012	9.752±0.013	10.463±0.012	0.728	0.711
BD+68 220	–	55858.6–55858.6	2	9.146±0.000	9.752±0.010	9.802±0.002	0.606	0.050
BD+43 1168	–	56002.3–56002.3	3	9.390±0.011	10.311±0.011	10.579±0.014	0.920	0.269

Table 6: Be stars observed at Hvar Observatory

Name of Star	HD/BD number	N of observations	Comparison HD/BD	Check HD/BD
10 Cas	144	18	2626	2011
V742 Cas	698	6	2626	2011
κ Cas	2905	17	2626	2011
ϕ Cas	4180	895	4142	6114
γ Cas	5394	137	2626	2011
V442 And	6226	964	4142	6114
φ And	6811	504	4142	6114
HD 9709	9709	7	4142	6114
φ Per	10516	678	12303	11291
V777 Cas	11606	11	12303	11291
V780 Cas	12302	9	12303	11291
HR 654	13854	13	12303	11291
V554 Per	14818	146	12303	11291
HR 894	18552	118	18411	19736
RX Cas	+67 244	361	18962	19556, 19193
HR 1051	21551	30	21278	24546
ψ Per	22192	29	21278	24546
HR 1113	22780	41	21856	23193
13 Tau	23016	114	23324	23288
ϕ Per	23180	33	21856	23193
17 Tau	23302	367	23324	23288
V971 Tau	23480	265	23324	23288
HR 1160	23551	17	21278	24546
η Tau	23630	205	23324	23288
BU Tau	23862	469	23324	23288
MX Per	25940	32	21278	24546
HR 1500	29866	30	33641	29722
BV Cam	32343	9	39283	34787
λ Eri	33328	24	32249	33224
V960 Tau	36576	392	36589	37711, 36819
zeta Tau	37202	1100	36589	37711, 36819
ω Ori	37490	90	36591	36351

Table 6: continued

Name of Star	HD/BD number	N of observations	Comparison HD/BD	Check HD/BD
V731 Tau	37967	107	36589	37711
V696 Mon	41335	318	42690	45546, 43023
HR 2231	43285	36	44783	43526
AX Mon	45910	40	44783	43526
HR 2370	45995	7	44783	43526
HD 46150	46150	3	44783	43526
HR 2418	47054	63	42690	45546, 43023
ψ 9 Aur	50658	29	49949	52860, 50860
AU Mon	50846	15	50109	50169
OT Gem	58050	437	58187	59059
β CMi	58715	168	58187	59059
BR CMi	61273	103	58187	61341
UX Mon	65607	152	65199	65005
HR 3135	65875	24	63975	71155
HD 81357	81357	93	82861	77692, 81772
κ Dra	109387	429	107193	115612, 104316
ϑ CrB	138749	158	138341	136849
V839 Her	142926	685	144206	145389, 141930
δ Sco	143275	98	144470	142096
χ Oph	148184	11	144470	142096
ζ Oph	149757	168	148367	147550
V2315 Oph	161261	5	161677, 169420	—
V744 Her	162732	1449	158414, 162132	162579
V2048 Oph	164284	164	164432	163641
V974 Her	164447	183	166182	166230
o Her	166014	182	166182	166230
NW Ser	168797	327	170200	169578
CX Dra	174237	1167	173664	172883
β Lyr	174638	544	176437	174602, 172044
V4024 Sgr	178175	1	177817	182678, 182645
ν Sgr	181616	46	177817	182678, 182645
HD 183261	183261	1	184606	188260
7 Vul	183537	131	188260	184606
V923 Aql	183656	1620	183227	184663
β 2 Cyg	183914	14	188260	184606
V1294 Aql	184279	1709	183227	184663
HR 7482	185859	38	188260	184606
V1507 Cyg	187399	193	188170	186357
V395 Vul	187811	39	188260	184606
V1746 Cyg	189687	209	188892	193369
V1624 Cyg	191610	510	188892	193369
20 Vul	192044	130	190993	191747
QR Vul	192685	191	190993	191747
P Cyg	193237	122	188892	193369
25 Vul	193911	124	190993	191747
V2119 Cyg	194335	140	188892	193369
V2120 Cyg	194883	74	194668	197618
HR 7843	195554	161	194668	197618
V1661 Cyg	198478	288	203245	199311
HR 7983	198625	78	203245	199311
V2140 Cyg	199478	90	203245	199311
V832 Cyg	200120	849	203245	199311
V1931 Cyg	200310	889	203245	199311
HR 8103	201733	71	203245	199311
ν Cyg	202904	6	202349	204403
HR 8153	203025	8	208218	202214
V382 Cep	203467	82	208218	202214
HR 8259	205551	21	207330	206259, 207793
ϵ Cap	205637	2	144206, 213420	—
HD 206773	206673	57	208218	202214

Table 6: continued

Name of Star	HD/BD number	N of observations	Comparison HD/BD	Check HD/BD
EM Cep	208392	10	208218	202214
HR 8375	208682	8	208218	202214
8 Lac A	214168	151	217101	214680
HR 8682	216057	205	218470	219080, 212593
V360 Lac	216200	424	217101	214680
EW Lac	217050	1281	218470	219080, 212593
V378 And	217543	267	217101	214680
σ And	217675	1636	217101	214680
KX And	218393	1210	218470	219080, 212593
KY And	218674	970	218470	219080, 212593
LQ And	224559	590	223229	222439, 224342
V639 Cas	225094	22	2626	2011
MWC 327	227611	6	+35 3955	190919
V1322 Cyg	229221	2	229234	229238

Table 7: Be stars observed at Hvar Observatory

Name of Star	JD range -2400000	V range (mag.)	B range (mag.)	U range (mag.)	B – V range (mag.)	U – B range (mag.)
10 Cas	45307.3–56515.6	5.558–5.614	5.534–5.582	5.332–5.385	-0.044 – -0.005	-0.223 – -0.196
V742 Cas	55791.5–55798.5	7.078–7.089	7.263–7.274	6.854–6.870	+0.178 – +0.187	-0.412 – -0.397
κ Cas	45307.3–56515.6	4.115–4.211	4.255–4.363	3.486–3.595	+0.129 – +0.163	-0.786 – -0.756
σ Cas	45212.6–60583.5	4.301–4.652	4.291–4.685	3.691–4.012	-0.073 – +0.033	-0.673 – -0.512
γ Cas	45307.3–60362.3	2.084–2.245	1.954–2.153	0.944–1.123	-0.141 – +0.005	-1.078 – -0.984
V442 And	45212.6–60583.5	6.572–6.897	6.592–6.867	5.966–6.327	-0.064 – +0.033	-0.646 – -0.540
φ And	45212.6–60583.5	4.217–4.308	4.163–4.271	3.801–3.932	-0.085 – -0.030	-0.403 – -0.324
HD 9709	53026.3–54356.5	7.085–7.179	7.022–7.155	6.586–6.740	-0.066 – -0.007	-0.436 – -0.414
φ Per	44935.4–60583.5	3.927–4.159	3.795–4.049	2.885–3.171	-0.188 – -0.036	-0.952 – -0.815
V777 Cas	53026.3–53388.3	6.902–7.040	6.976–7.096	6.152–6.286	+0.044 – +0.076	-0.837 – -0.802
V780 Cas	53028.3–53388.3	8.042–8.102	8.322–8.374	7.842–7.936	+0.268 – +0.284	-0.482 – -0.435
HR 654	44985.3–46696.6	6.446–7.383	6.734–7.621	6.094–6.995	+0.234 – +0.292	-0.644 – -0.615
V554 Per	44935.4–50087.2	6.187–6.318	6.486–6.638	5.878–6.054	+0.282 – +0.327	-0.618 – -0.545
HR 894	44860.6–54026.6	6.105–6.181	6.036–6.108	5.642–5.710	-0.096 – -0.024	-0.424 – -0.372
RX Cas	42603.5–60330.3	8.646–9.500	9.777–11.037	9.959–12.169	+0.919 – +1.554	-0.114 – +1.132
HR 1051	45307.4–46689.6	5.828–5.871	5.787–5.814	5.469–5.495	-0.059 – -0.027	-0.327 – -0.302
ψ Per	45307.4–46689.6	4.194–4.247	4.106–4.136	3.508–3.547	-0.135 – -0.073	-0.604 – -0.573
HR 1113	44988.3–50087.5	5.551–5.596	5.491–5.524	5.060–5.104	-0.088 – -0.048	-0.440 – -0.409
13 Tau	44863.6–54015.5	5.668–5.710	5.660–5.703	5.371–5.452	-0.029 – +0.009	-0.302 – -0.220
σ Per	44988.3–45359.3	3.804–3.853	3.853–3.910	3.128–3.177	+0.024 – +0.077	-0.748 – -0.702
17 Tau	44863.6–60373.3	3.638–3.764	3.539–3.672	3.149–3.298	-0.152 – -0.073	-0.455 – -0.345
V971 Tau	44863.6–60373.3	4.170–4.235	4.119–4.187	3.679–3.816	-0.090 – -0.027	-0.471 – -0.344
HR 1160	45307.4–46689.6	6.171–6.200	6.236–6.257	5.915–5.938	+0.043 – +0.070	-0.327 – -0.303
η Tau	44863.6–60373.3	2.875–2.962	2.735–2.854	2.385–2.505	-0.158 – -0.062	-0.393 – -0.303
BU Tau	43140.3–60583.5	4.999–5.405	4.919–5.377	4.452–5.241	-0.113 – -0.001	-0.476 – +0.019
MX Per	43136.4–46689.6	3.984–4.075	3.965–4.046	3.408–3.472	-0.106 – +0.014	-0.574 – -0.547
HR 1500	44921.5–52202.6	6.058–6.096	6.126–6.158	5.813–5.871	+0.046 – +0.086	-0.334 – -0.275
BV Cam	45308.6–45697.4	5.139–5.184	5.077–5.099	4.336–4.354	-0.089 – -0.060	-0.748 – -0.737
λ Eri	51445.6–51520.5	4.246–4.298	4.060–4.117	3.172–3.215	-0.217 – -0.137	-0.925 – -0.871
V960 Tau	44902.6–60583.6	5.368–5.718	5.447–5.737	4.690–4.980	-0.016 – +0.136	-0.832 – -0.719
ζ Tau	43135.4–60583.6	2.769–3.200	2.570–2.993	1.757–2.369	-0.278 – -0.087	-0.866 – -0.614
ω Ori	51512.5–59598.4	4.441–4.597	4.371–4.485	3.601–3.726	-0.134 – -0.070	-0.776 – -0.732
V731 Tau	43136.4–46694.6	6.197–6.270	6.141–6.199	5.489–5.583	-0.114 – -0.034	-0.673 – -0.603
V696 Mon	44902.6–59974.4	5.179–5.302	5.117–5.232	4.280–4.396	-0.092 – -0.015	-0.876 – -0.780
HR 2231	44993.4–50865.3	6.058–6.133	5.943–6.037	5.411–5.529	-0.140 – -0.099	-0.532 – -0.479
AX Mon	44993.4–53388.5	6.684–6.810	6.988–7.134	6.344–6.771	+0.282 – +0.380	-0.661 – -0.351
HR 2370	49730.4–49751.4	6.058–6.101	6.000–6.031	5.152–5.195	-0.078 – -0.058	-0.848 – -0.820
HD 46150	53388.5–53388.5	6.716–6.728	6.875–6.868	6.048–6.051	+0.132 – +0.145	-0.817 – -0.807
HR 2418	44902.6–52561.6	5.501–5.554	5.411–5.466	5.022–5.099	-0.107 – -0.055	-0.411 – -0.340
ψ 9 Aur	45014.4–52940.5	5.833–5.893	5.769–5.828	5.325–5.391	-0.096 – -0.054	-0.454 – -0.419

Table 7: continued

Name of Star	JD range -2400000	V range (mag.)	B range (mag.)	U range (mag.)	B – V range (mag.)	U – B range (mag.)
AU Mon	54862.4–55104.6	8.221–9.134	8.275–9.323	7.648–8.804	-0.003 – +0.193	-0.637 – -0.481
OT Gem	44977.4–58924.4	6.061–6.509	5.944–6.346	5.097–5.541	-0.230 – -0.071	-0.922 – -0.760
β CMi	44977.4–58564.4	2.862–2.917	2.755–2.844	2.473–2.593	-0.128 – -0.062	-0.308 – -0.242
BR CMi	55574.5–56015.3	7.089–7.184	7.355–7.433	7.392–7.483	+0.233 – +0.299	+0.005 – +0.076
UX Mon	52655.4–55594.4	8.230–9.343	8.569–10.112	8.341–10.789	+0.286 – +0.957	-0.233 – -0.717
HR 3135	45056.3–53035.5	6.444–6.505	6.348–6.428	5.602–5.653	-0.133 – -0.052	-0.791 – -0.721
HD 81357	55879.6–57116.4	8.295–8.391	8.459–8.553	8.264–8.334	+0.118 – +0.186	-0.251 – -0.172
κ Dra	42237.4–60454.4	3.766–3.978	3.624–3.882	3.071–3.353	-0.218 – +0.049	-0.637 – -0.483
θ CrB	44801.4–58637.5	4.128–4.230	3.986–4.138	3.443–3.628	-0.164 – -0.073	-0.570 – -0.449
V839 Her	41528.5–60520.4	5.705–5.803	5.572–5.710	5.158–5.352	-0.162 – -0.058	-0.447 – -0.291
δ Sco	52068.4–60518.4	1.579–2.231	1.630–2.330	0.705–1.208	-0.137 – -0.147	-1.130 – -0.861
χ Oph	54275.4–54290.4	4.328–4.427	4.637–4.737	3.896–3.970	+0.290 – +0.350	-0.781 – -0.732
ζ Oph	52494.3–58682.4	2.510–2.584	2.522–2.615	1.677–1.782	-0.027 – +0.071	-0.886 – -0.803
V2315 Oph	41896.4–42625.4	8.240–8.461	8.275–8.552	8.106–8.398	+0.035 – +0.100	-0.198 – -0.105
V744 Her	41535.4–60580.3	6.662–6.945	6.504–6.858	6.002–6.519	-0.191 – -0.058	-0.557 – -0.265
V2048 Oph	45105.5–60545.4	4.636–4.898	4.616–4.787	3.737–4.056	-0.132 – -0.001	-0.889 – -0.703
V974 Her	45065.6–58395.3	6.405–6.586	6.339–6.508	5.928–6.134	-0.101 – -0.003	-0.419 – -0.321
σ Her	45065.6–58395.3	3.784–3.892	3.720–3.921	3.669–3.851	-0.123 – +0.045	-0.142 – +0.033
NW Ser	44433.4–48133.3	6.019–6.191	6.042–6.173	5.381–5.595	-0.041 – +0.035	-0.666 – -0.578
CX Dra	42977.3–56094.4	5.666–6.014	5.658–5.959	4.890–5.252	-0.143 – +0.025	-0.818 – -0.642
β Lyr	49272.3–57949.5	3.324–4.269	3.303–4.323	2.745–3.803	-0.052 – -0.090	-0.628 – -0.449
V4024 Sgr	53584.4–53584.4	5.309–5.309	5.308–5.308	4.544–4.544	-0.001 – -0.001	-0.764 – -0.764
ν Sgr	53570.4–54297.4	4.510–4.620	4.627–4.769	4.055–4.190	+0.093 – +0.149	-0.597 – -0.555
HD 183261	54275.5–54275.5	6.865–6.865	6.854–6.854	6.167–6.167	-0.011 – -0.011	-0.687 – -0.687
7 Vul	54273.4–59025.5	6.317–6.388	6.219–6.306	5.677–5.771	-0.109 – -0.073	-0.561 – -0.503
V923 Aql	44073.4–60580.3	5.983–6.166	5.966–6.201	5.618–6.018	-0.044 – +0.095	-0.391 – -0.102
β_2 Cyg	46679.4–46695.4	5.140–5.159	5.051–5.069	4.730–4.744	-0.098 – -0.078	-0.327 – -0.315
V1294 Aql	44073.4–60580.4	6.788–7.565	6.807–7.634	5.997–7.587	-0.005 – +0.173	-0.841 – +0.002
HR 7482	46679.4–58682.5	6.481–6.519	6.859–6.913	6.270–6.326	+0.376 – +0.403	-0.613 – -0.577
V1507 Cyg	43024.4–43760.4	6.896–7.088	7.082–7.260	6.680–6.861	+0.103 – +0.263	-0.463 – -0.323
V395 Vul	46679.4–58682.5	4.909–4.957	4.726–4.792	4.067–4.137	-0.183 – -0.141	-0.700 – -0.644
V1746 Cyg	46245.5–60583.3	5.121–5.246	4.990–5.084	4.251–4.409	-0.168 – -0.108	-0.747 – -0.670
V1624 Cyg	42237.5–60583.3	4.888–5.084	4.721–4.943	3.943–4.281	-0.216 – -0.103	-0.850 – -0.639
20 Vul	44432.5–60565.5	5.867–5.945	5.746–5.842	5.285–5.405	-0.135 – -0.089	-0.485 – -0.428
QR Vul	44794.5–60583.3	4.600–4.820	4.453–4.644	3.684–3.925	-0.227 – -0.109	-0.799 – -0.706
P Cyg	46287.4–55451.4	4.715–4.877	5.116–5.275	4.559–4.718	-0.383 – -0.450	-0.581 – -0.513
25 Vul	44817.4–60565.5	5.499–5.563	5.395–5.487	4.958–5.078	-0.120 – -0.073	-0.460 – -0.387
V2119 Cyg	46238.5–47022.4	5.783–5.923	5.616–5.758	4.652–4.836	-0.190 – -0.141	-0.981 – -0.906
V2120 Cyg	45149.5–55068.4	7.163–7.361	7.221–7.345	6.551–6.730	-0.019 – +0.060	-0.672 – -0.609
HR 7843	44431.5–55064.4	5.863–5.942	5.827–5.880	5.574–5.675	-0.095 – -0.009	-0.253 – -0.166
V1661 Cyg	41528.4–53935.5	4.740–4.896	5.112–5.272	4.651–4.835	+0.333 – +0.424	-0.472 – -0.404
HR 7983	46696.4–53592.5	6.323–6.400	6.264–6.342	5.674–5.796	-0.074 – -0.038	-6.616 – -0.546
V2140 Cyg	43024.4–54662.5	5.634–5.724	6.097–6.192	5.787–5.900	+0.424 – +0.486	-0.334 – -0.283
V832 Cyg	46258.6–60583.4	4.451–4.837	4.432–4.782	3.489–3.887	-0.116 – +0.013	-0.984 – -0.860
V1931 Cyg	46281.5–60583.4	5.289–5.503	5.136–5.303	4.152–4.405	-0.251 – -0.122	-1.001 – -0.890
HR 8103	46696.4–52100.5	6.595–6.650	6.454–6.514	5.747–5.890	-0.164 – -0.124	-0.707 – -0.621
ν Cyg	44817.4–44818.5	4.342–4.371	4.256–4.284	3.433–3.460	-0.094 – -0.081	-0.828 – -0.815
HR 8153	48116.6–48123.6	6.378–6.443	6.585–6.666	6.085–6.220	+0.191 – +0.227	-0.507 – -0.446
V382 Cep	45536.4–52282.2	4.924–5.255	4.942–5.222	4.299–4.651	-0.128 – +0.051	-0.694 – -0.527
HR 8259	45155.5–59078.6	6.153–6.202	6.139–6.199	5.875–5.947	-0.026 – +0.001	-0.283 – -0.238
ϵ Cap	41929.5–41940.4	4.582–4.613	4.453–4.534	3.811–3.880	-0.128 – -0.079	-0.654 – -0.642
HD 206773	52142.6–52282.2	6.876–7.053	6.980–7.183	6.169–6.428	+0.087 – +0.143	-0.830 – -0.742
EM Cep	41954.5–54020.3	6.975–7.115	7.224–7.398	6.665–6.851	+0.239 – +0.296	-0.579 – -0.540
HR 8375	48116.6–48123.6	5.896–5.939	5.807–7.873	5.062–5.131	-0.102 – -0.055	-0.747 – -0.715
8 Lac A	43013.6–54026.5	5.640–5.767	5.483–5.662	4.575–4.723	-0.191 – -0.064	-0.950 – -0.853
HR 8682	44432.6–51426.5	6.117–6.199	6.030–6.114	5.528–5.674	-0.125 – -0.043	-0.536 – -0.440
V360 Lac	44473.5–53980.4	5.862–5.994	5.949–6.067	5.442–5.579	+0.041 – +0.112	-0.554 – -0.469
EW Lac	41534.5–60199.5	5.096–5.492	4.958–5.361	4.198–4.955	-0.165 – -0.030	-0.778 – -0.397
V378 And	42261.5–55460.4	6.461–6.615	6.317–6.530	5.630–5.906	-0.180 – -0.061	-0.743 – -0.624
σ And	42614.6–60583.4	3.498–3.784	3.395–3.702	2.861–3.256	-0.178 – +0.000	-0.598 – -0.427

Table 7: continued

Name of Star	JD range -2400000	<i>V</i> range (mag.)	<i>B</i> range (mag.)	<i>U</i> range (mag.)	<i>B</i> – <i>V</i> range (mag.)	<i>U</i> – <i>B</i> range (mag.)
KX And	42261.5–57621.5	6.775–7.212	7.092–7.596	6.661–7.478	+0.251 – +0.431	-0.551 – -0.101
KY And	42614.5–57621.5	6.686–6.894	6.679–6.862	6.075–6.317	-0.092 – +0.059	-0.642 – -0.513
LQ And	44433.5–60583.4	6.498–6.598	6.412–6.524	5.793–5.920	-0.125 – -0.035	-0.669 – -0.554
V639 Cas	45307.3–55798.5	6.201–6.273	6.542–6.610	6.011–6.099	+0.326 – +0.347	-0.532 – -0.506
MWC 327	44841.4–46689.4	8.710–8.790	9.074–9.148	8.384–8.435	+0.335 – +0.364	-0.721 – -0.666
V1322 Cyg	44794.5–44843.3	9.333–9.346	10.190–10.223	9.957–10.016	+0.844 – +0.890	-0.233 – -0.207